

# Age Estimation using Pulp Chamber/Crown Volume Ratio of Permanent Maxillary and Mandibular Second Molars



Rasha Ibrahim Alghali<sup>1</sup>, Phrabhakaran Nambiar<sup>2</sup>, Wei Cheong Ngeow<sup>1,\*</sup>, Norliza Ibrahim<sup>1</sup> and Muhammad Khan Asif<sup>2</sup>

<sup>1</sup>Department of Oral & Maxillofacial Clinical Sciences, Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia

<sup>2</sup>Department of Preclinical Sciences, MAHSA University, Faculty of Dentistry, Bandar Saujana Putra, Malaysia

## Abstract:

**Introduction/Objective:** Age estimation is an essential component in defining multiple elements of the biological profile of unidentified individuals or people with undocumented age. This research examines the relationship between chronological age and the ratio of pulp chamber volume to crown volume in maxillary and mandibular second molars within a Malaysian population (comprising Malays and Chinese).

**Methods:** The study employs volumetric analysis using CBCT scans, enhanced with Mimics software, to evaluate this association. A total of 316 CBCT scans from 154 Malays and 162 Chinese, aged 18 to 65 years, were categorized into four distinct age groups for analysis.

**Results:** In this study, Pearson correlation analysis showed that the mandibular second molar (LM2) (-0.746) exhibited a slightly greater coefficient of correlation (R) compared to the maxillary second molar (UM2) (-0.739). The Fisher Z test demonstrated that dental age estimation is sex-independent. The study's newly derived regression equation was validated on a separate set of 68 permanent second molars. The findings revealed Mean Absolute Error (MAE) results of 5.3 and 7.5 years for maxillary and mandibular second molars, respectively.

**Discussion:** The PC/CVR model showed a significant inverse correlation with age and higher accuracy in individuals under 50 years; however, performance decreased in older age groups (50+).

**Conclusion:** This study found that a volumetric change in the pulp chamber with age is a valid method for estimating dental age in the Malaysian population.

**Keywords:** Dental age estimation, CBCT, Mimics software, Malaysian population, Second molars.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

\*Address correspondence to this author at the Department of Oral & Maxillofacial Clinical Sciences, Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia; E-mail: [ngeow@um.edu.my](mailto:ngeow@um.edu.my)

Cite as: Alghali R, Nambiar P, Ngeow W, Ibrahim N, Asif M. Age Estimation using Pulp Chamber/Crown Volume Ratio of Permanent Maxillary and Mandibular Second Molars. Open Dent J, 2026; 20: e187421062306. <http://dx.doi.org/10.2174/01187421062306260924075306>



CrossMark

Received: December 24, 2025

Revised: June 24, 2026

Accepted: July 27, 2026

Published: October 01, 2026



Send Orders for Reprints to [reprints@benthamscience.net](mailto:reprints@benthamscience.net)

## 1. INTRODUCTION

Age estimation is an important component of forensic medicine and is applied in accidents, criminal investigations, and civil matters (hiring, marriage, and school attendance) [1]. Undocumented children have a

higher risk of having their rights violated. Age estimation can also be applied to cases of missing documentation of adopted children and illegal immigrants [2, 3].

Dental age estimation is one of the most valid approaches of age determination, especially in individuals

aged less than 25 years [4]. Teeth are made of calcified tissues, which are more resistant to mechanical forces, harsh chemical conditions, and extreme heat than other tissues. As a result, they can survive extreme conditions, particularly the upper/maxillary second molars (UM2) and lower/mandibular second molars (LM2). Posterior teeth are generally more resistant to the same hazardous conditions than the anterior teeth, as they are naturally shielded by the surrounding tissues, "i.e., the cheeks and the tongue". As a result, human teeth, especially posterior teeth, are considered to be important biological tissues for human identification [5].

Techniques developed to estimate dental age using developmental phases and the eruption pattern have made age estimation easier among children and adolescents [6, 7]. In childhood, dental age can be determined using tooth development charts, whereas in young adulthood, crown and root morphology of the third molar is used to estimate age. Different stages of mineralization of permanent teeth are highly beneficial until the age of 15-16 years, which is when the development of the second molars is completed. Secondary dentin deposition is a reliable physiological age-related parameter that can be investigated to estimate an adult's age [8, 9]. Physiological changes in teeth and their surrounding structures occur throughout a human's life. Once morphological changes in dental tissues are complete at age 25 (when the third molar is fully formed), examining the diminution of the pulp cavity resulting from secondary dentin formation becomes a useful physiological parameter for age estimation. Three-dimensional (3D) cone-beam computed tomography (CBCT) enables assessment of oral tissues with limited X-ray radiation exposure for patients [10]. This approach is one of the most reliable [7, 11-13] and least invasive methods for age determination among adults, as no tissue needs to be extracted from the body [14]; thus, this method can be applied to both living and deceased individuals. Cameriere and Ferrante [15] initially tested the pulp/tooth ratio method on periapical radiographs. Several researchers have since validated it on 3D scans [11, 16-20]. Currently, this approach is widely recognized and applied worldwide as a reliable, less intrusive technique for determining adult age. The original Cameriere approach was applied to many populations using CBCT scans [8, 11, 21, 22].

In previous studies, Asif and Nambiar [23] and Akbarizadeh and Tabrizi [24] analyzed CBCT images and found that the pulp chamber/crown (PC/C) approach had a stronger association with age than the pulp/tooth (P/T) approach. Realizing the potential of the pulp chamber/crown approach for age estimation, the authors adopted this method in the present study.

Despite numerous studies demonstrating that ethnic background and population origin significantly influence 3D dental pulp cavity volume and volumetric ratios [8, 25], other studies reported no difference regarding age estimation between ethnicities [26, 27]. However, various studies have suggested that a population-specific model remains essential for accurate age estimation.

The objective of this research was to formulate a

regression equation for estimating age through studying the strength of correlation between the chronological age and the PC/C volume ratio of permanent second molars in a selected Malaysian population. This was performed using 3D CBCT scans of the permanent UM2 and LM2, as these teeth have greater longevity than the first molars and are less variable in morphology than the third molars. Statistical differences in the strength of correlation between sex (male and female) and ethnicity (Malays and Chinese) were also investigated. Subsequently, the new regression equation was applied to independent CBCT images to validate it.

## 2. MATERIALS AND METHODS

The ethical approval for this retrospective study was obtained from the University of Malaya Medical Ethics Committee (Ref. No: DF OS2002/0003(L)). This study was conducted in accordance with the ethical principles of the Declaration of Helsinki [28]. Patients were well aware that the data from their CBCT scans could be utilized for research and education at the faculty, and consented to their use. CBCT scans of 316 patients, 154 Malays and 162 Chinese, were retrieved from the database stored in the Oral and Maxillofacial Imaging Division, Faculty of Dentistry, University of Malaya, with exposure parameters of 120 KV and 18 mA. All scans were obtained with a medium Field Of View (FOV), a voxel size of 0.30 mm, and a scanning period of 20 seconds. The images were selected based on certain inclusion criteria, as only high-quality CBCT images of Malay and Chinese Malaysians aged 18 to 65 years with no evidence of caries or pathology associated with the targeted tooth were selected. Poor-quality CBCT images, or those not acquired using the scanning parameters of this study, and patients outside the inclusion age were excluded. In addition, any restorations, caries, or pathology associated with targeted teeth, as well as teeth with obliterated cemento-enamel junctions or open apices, were excluded from the study. Sex and Gender Equity in Research (SAGER) guidelines were followed for reporting sex in this study [29].

To maintain a balanced and equally distributed sample, patients whose age ranged from 18 to 65 years were selected. The data were then categorized into four groups with 11-year intervals except for the above 50-year-old group.

The sample size of the study was calculated using G\*Power software (version 3.1.9.4) by employing an F-test, Power of  $(1 - \beta \text{ err})$  0.95, and  $\alpha=0.05$ . The effect size (0.15) was determined by computing a 0.131 coefficient of determination ( $R^2$ ) value. The calculated sample size was 137; however, 210 UM2 and 197 LM2 were included in this study.

All selected CBCT scans were stored in Digital Imaging and Communications in Medicine (DICOM) format. A total of 316 high-quality scans stored from the i-CAT Cone Beam 3D Dental Imaging System (version 3.1.62 supplied by Imaging Sciences International, Hatfield, USA) were selected based on the patients' age and ethnicity. Subsequently, 407 fully developed non-carious intact teeth

(210 UM2 and 197 LM2) were retrieved from the database. Only one UM2 and one LM2 were selected per image to prevent any biased results when data were obtained twice from the same subject. Besides the study sample of 407 teeth, an independent sample of 68 UM2 and LM2 was selected to validate the newly developed regression equation.

## 2.1. Pulp chamber-to-crown Volume Ratio (PC/CVR) Measurement According to Asif *et al.* is as follows [8]

- [1] The images were transferred to the Mimics software (Materialise NV, Belgium, version 16.0) as DICOM files.
- [2] The CBCT scans were properly positioned in the axial, coronal, and sagittal views, so the targeted tooth could be identified in all three views.
- [3] Using the segment tab, various thresholds were established to generate a binary mask that retained voxels within the upper and lower ranges for different parts (pulp and tooth).
- [4] To isolate the tooth from adjacent structures, masks were meticulously reviewed segment by segment during the multiple-slice editing stage to ensure accurate segmentation and separation.
- [5] The examiner chose a specific seed point to start a region growing within the targeted structure of the selected tooth, and new masks were generated for the targeted parts (pulp and tooth) of the investigated tooth, excluding unwanted surrounding structures.
- [6] Specific refining tools, such as 'Edit Masks' and 'Multiple Slice Edit', were used for the manual inclusion or removal of unnecessary voxels.
- [7] Following the mask editing process, the masks of the targeted pulp and tooth were selected, and then the 'Calculate Part' function was utilized to generate 3D-reconstructed models of both the pulp tissue and the tooth (Fig. 1).
- [8] After generating three-dimensional models of the targeted pulp and tooth, a plane was established by selecting three points on the Cementoenamel Junction (CEJ) in the sagittal view. This plane was then aligned with the CEJ of the targeted tooth in both the coronal and axial views. Following the methodology described by Azim and Azim [30] for UM2, the floor of the pulp chamber in the coronal view was used as a reference point to distinguish the anatomical crown from the root. For LM2, the sagittal view was utilized, with the pulp chamber floor serving as a reference to separate the crown from the root [31].
- [9] The plane on the 3D model was adjusted to align with the pulp chamber floor. Subsequently, the crown and root were separated using 3D tools at the level of the plane (Fig. 2). After selecting 'Properties' from the menu, the volume value ( $\text{mm}^3$ ) displayed in the box was recorded for every 3D reconstructed pulp chamber and each 3D reconstructed crown model.
- [10] Using Excel 2016, Pulp Chamber Volume (PCV) was divided by Crown Volume (CV) to obtain the Pulp Chamber/Crown Volume Ratio (PC/CVR).





Fig. (1). Pulp and tooth thresholding and the new mask of the maxillary left second molar.



**Fig. (2).** **A:** The plane separating different parts of a maxillary second molar 3D model. **B:** The separated pulp chamber and crown on the plane.

**3. RESULTS**

To ensure the reproducibility of this method, an intraclass correlation coefficient (ICC) analysis was conducted to assess both intra-examiner and inter-examiner reliability. Volumetric measurements were performed on twenty randomly selected teeth, comprising ten UM2 and ten LM2, over a three-month period to evaluate intra-examiner reliability. Additionally, a second examiner, a postgraduate dental student, performed a volumetric analysis on the same sample to determine inter-examiner reliability. The ICC values obtained were 0.973 for intra-examiner reliability and 0.981 for inter-examiner reliability; these values are classified as

excellent. These results demonstrate that the method is highly consistent and reproducible.

A statistical analysis using Fisher z-test showed no significant difference between ethnicity (Table 1) and sex (Table 2).

In summary, there was no significant difference in the pulp chamber/crown volume ratio (PC/CVR) of UM2 and LM2 when compared by sex and ethnicity. As a result, the PC/CVR was deemed more appropriate for calculation across the entire study sample, examining the relationship between the PC/CVR of UM2 and LM2 and chronological age. This relationship was found to be significantly inverse ( $p < 0.05$ ) (Table 3).

**Table 1.** Fisher’s Z test for the investigated teeth among males and females, based on ethnicities.

| Variables |     | r      |         | Fisher’s Z test |         |
|-----------|-----|--------|---------|-----------------|---------|
|           |     | Malay  | Chinese | Z-value         | P-value |
| Males     | UM2 | -0.752 | -0.759  | 0.082           | 0.467   |
|           | LM2 | -0.738 | -0.755  | 0.186           | 0.426   |
| Females   | UM2 | -0.727 | -0.791  | 0.743           | 0.229   |
|           | LM2 | -0.772 | -0.759  | -0.149          | 0.441   |

**Table 2.** Fisher’s Z test for the investigated teeth among Malays and Chinese, based on sex.

| Variables |     | r      |        | Fisher’s Z test |         |
|-----------|-----|--------|--------|-----------------|---------|
|           |     | Male   | Female | Z-value         | P-value |
| Malays    | UM2 | -0.776 | -0.727 | -0.272          | 0.393   |
|           | LM2 | -0.738 | -0.772 | 0.365           | 0.358   |
| Chinese   | UM2 | -0.759 | -0.791 | 0.402           | 0.344   |
|           | LM2 | -0.755 | -0.759 | 0.047           | 0.481   |

**Table 3. The correlation coefficient between chronological age and pulp chamber/crown volume ratio (PC/CVR) for UM2 and LM2 for the whole sample.**

| Investigated Variable | UM2 (r)  | LM2 (r)  | UM2 and LM2 (r) |
|-----------------------|----------|----------|-----------------|
| pulp chamber/crown VR | -0.739** | -0.746** | -0.741*         |

Note: \*\* Significant at the 0.01 probability level

\* Significant at the 0.05 probability level

An independent *t*-test was applied to compare the effect of age within the two ethnicities. There was no significant difference in the mean values of PC/C volume ratio between ethnicities in the three age groups. However, the 29-39 age group showed a significant difference between Malay and Chinese populations (Table 4).

A multiple linear regression analysis was conducted to generate a newly derived age estimation model for Malaysian adults. A stepwise regression analysis was used to identify predictor variables that significantly contributed to the model. This is particularly useful in the current case as many predicted variables are included, so the non-significant predictors are excluded successively. It also shows different potential models, and therefore allows the researchers to choose the best predictive model. The results of the study indicated a significant linear relationship between the dependent and the independent variables. The results also showed that the assumptions of no multicollinearity and normal distribution of residuals were met.

The model yielded an R-square ( $R^2$ ) value of 0.543 and an adjusted R-square value of 0.540, indicating that 54.3% of the variation in age could be explained by the PC/CVR. Using stepwise regression analysis to derive an age estimation regression model, a forward selection approach was applied. All possible predictors were included in the

model; gradually, the non-significant predictors were successively removed until the best model was built. Using stepwise regression analysis indicated that the *p*-value was less than 0.005 for ethnicity and sex, indicating that ethnicity has a significant effect on the estimated age (Table 5). Therefore, ethnicity and sex were included in the model. However, tooth type was excluded as the *p*-value (0.239) and standardized coefficient (-0.040) showed that it was not contributing significantly to the regression formula.

Fisher z-test showed a significant difference between the study and validation samples with a *p*-value of 0.014 (Table 6). However, both groups exhibited a strong correlation with age.

The age estimation equation derived from the study was then evaluated using an independent validation group, consisting of 38 UM2 and 30 LM2. The results demonstrated acceptable values for the Mean Absolute Error (MAE) for the pulp chamber/crown volume ratio (PC/C VR) in both groups of teeth. This indicates that the derived age estimation formula is reliable and can serve as a valid method for estimating age among Malaysian adults. However, it was observed that all variables exhibited the weakest correlation in the age group above 50 years (Table 7).

**Table 4. Independent t-test analysis of PC/CVR mean values for the whole study sample for different age groups based on ethnicities**

| Age Group | Malay Mean (SD) | Chinese Mean (SD) | t test  | p-value |
|-----------|-----------------|-------------------|---------|---------|
| 18-28     | 0.0728 (0.018)  | 0.0740 (0.013)    | 0.6819  | 0.495   |
| 29-39     | 0.0492 (0.011)  | 0.0621 (0.011)    | 10.4201 | 0.000   |
| 40-50     | 0.0460 (0.011)  | 0.0486 (0.010)    | 2.2004  | 0.028   |
| 50+       | 0.0391 (0.011)  | 0.0405 (0.009)    | 1.2409  | 0.215   |

**Table 5. Stepwise regression analysis to derive an age estimation regression model.**

| Model        | Value    | Std. Error | Standardized Coefficient | t- value | P - value | 95% Confidence Interval |             | VIF   |
|--------------|----------|------------|--------------------------|----------|-----------|-------------------------|-------------|-------|
|              |          |            | Beta                     |          |           | Lower Bound             | Upper Bound |       |
| Intercept    | 66.643   | 2.469      |                          | 26.996   | 0.000     | 60.987                  | 69.090      |       |
| ethnicity    | 3.021    | 0.921      | 0.110                    | 3.281    | 0.001     | 1.178                   | 4.798       | 1.003 |
| sex          | 1.856    | 0.924      | -0.068                   | 2.008    | 0.045     | -3.678                  | -0.042      | 1.013 |
| measurements | -544.115 | 24.755     | -0.742                   | -21.980  | 0.000     | -591.942                | -494.606    | 1.016 |

Note: The derived regression formula for permanent second molars is as follows:

Age=66.643+3.021E-1.856S-544.115M

E= ethnicity (1=Malay, 2=Chinese) (categorical data)

S= sex (0=male, 1=female) (categorical data)

M= the obtained measurement of PC/CVR (continuous data)

**Table 6. The correlation coefficient between chronological age and pulp chamber/crown volume ratio (PC/CVR) for the validation and the study group.**

| Investigated Variable | Validation Group (r) | Study Group (r) | Z-value | P-value |
|-----------------------|----------------------|-----------------|---------|---------|
| pulp chamber/crown VR | -0.847**             | -0.741*         | 2.191   | 0.014   |

**Table 7. Mean Absolute Error (MAE) for different age groups.**

| Age Group  | Mean Absolute Error (MAE) (Year) |      |
|------------|----------------------------------|------|
|            | UM2                              | LM2  |
| 18-28      | 4.5                              | 6.5  |
| 29-39      | 3.9                              | 4.5  |
| 40-50      | 3.9                              | 4.2  |
| Above 50   | 9.1                              | 14.9 |
| All groups | 5.3                              | 7.5  |

4. DISCUSSION

Dental age estimation is one of the most reliable and valid methods of age estimation and identification in forensic sciences [23]. It is also considered valuable for identifying deceased individuals and determining the chronological age of living persons [6]. The disadvantage of using CBCT imaging is that it has limited visualization of the soft tissue. However, soft tissues are not important in some age estimation techniques [32]. Compared to other age-related physiological changes, tooth maturation exhibits less variability and has a strong correlation with chronological age [33]. Traditionally, age estimation relied on conventional imaging techniques that converted three-dimensional (3D) structures into two-dimensional (2D) images. This approach limited the visualization of crucial dental and surrounding tissue features, as they were primarily observed in the mesiodistal plane, while details in the buccolingual dimension were not fully captured [34]. Given the constraints associated with two-dimensional images, researchers have advocated for the use of 3D image analysis, specifically utilizing CBCT imaging to achieve more efficient diagnostic and treatment planning [35]. In addition, 3D imaging techniques have contributed to the age estimation domain with promising results, offering improved age estimation accuracy over traditional 2D radiography [8].

The deposition of secondary dentin occurs only after root development is completed [36]. As a result, teeth with open apices were not considered in this study. Given that secondary dentin formation may not yet have commenced in individuals under 18 years of age, they were also excluded. Consequently, the P/T (Pulp/Tooth ratio) is not a reliable method for age estimation in children and adolescents, and was not adopted [37, 38].

Previous studies reported a moderate correlation between chronological age and pulp volume [39-41]. However, pulp chamber volume as an indicator for age estimation remains debatable, as it varies between individuals, and even those of the same age may exhibit

differences in pulp size [42, 43]. In contrast, the PC/CVR employed in this study mitigates individual variations in pulp chamber size more effectively than relying solely on pulp chamber volume [22]. Moreover, using pulp chamber cavity volume as the numerator is recommended to avoid the issue of a zero-value denominator in cases of pulp chamber obliteration [44].

Advances in three-dimensional (3D) imaging and image-processing software have improved forensic age estimation by enabling precise volumetric evaluation of the dental pulp cavity [45], with some researchers having chosen to reassess traditional methodologies using these innovative 3D techniques. For instance, several studies have applied the Cameriere method, which involves calculating the P/T area ratio, through 3D imaging [46-48]. Their findings demonstrated a moderate to strong correlation between the P/T area ratio and chronological age. Specifically, Salemi *et al.* [48] reported correlation coefficients of  $-0.767$  for males and  $-0.794$  for females. Similarly, Sue *et al.* [46] examined the P/T area ratio in maxillary and mandibular first molars, yielding determination coefficients of  $R^2 = 0.586$  and  $R^2 = 0.609$ , respectively. Notably, these values align closely with the results obtained from our volumetric analysis, highlighting the consistency and reliability of 3D-based approaches in age estimation.

CBCT scans were used to assess the original Cameriere's approach on several populations. The current study's findings are compared to those of earlier research using CBCT pictures in this section. Yang *et al.* [21], Star *et al.* [22], and De Angelis *et al.* [18] reported a weak to moderate connection ( $R^2 < 0.52$ ) between the pulp/tooth volume ratio in single-rooted teeth and chronological age in the Belgian population. The PC/CVR determination coefficient values in our research ( $R^2 = 0.54$ ) are comparable to those found in their study.

Additionally, the correlation strength observed for the pulp chamber-to-crown volume ratio in second molars in this study was notably stronger ( $R = -0.741$ ) compared to

the correlations reported in previous research. Specifically, Aboshi and Takahashi [49], Yang and Fan [50], and Türk and Görmez [51] reported weaker correlations ( $R \geq -0.45$ ).

In other studies investigating pulp/tooth volume ratio, Zhan and Chen [52], Asif *et al.* [23], Krishnapriya and Debta [53], and Prakash [54] reported higher determination coefficients ( $R^2 > 0.625$ ) with age than those reported in the present study. However, it is worth mentioning that both studies investigated anterior teeth.

In this study, the association between age and the PC/CVR was stronger in males than in females for the UM2 among Malays. However, for the LM2 in Malays, females exhibited stronger correlation coefficient values than males. In contrast, among the Chinese population, females showed a stronger correlation than males for both UM2 and LM2. Notably, the Fisher's Z test results indicated no statistically significant differences in the correlation coefficients between sexes, aligning with findings from previous studies [11, 18, 23, 55-57].

Asif *et al.* [23] and De Micco *et al.* [25] emphasized the necessity of developing population-specific regression equations to accurately estimate dental age in adult populations. Studies comparing equations derived from external populations with those specifically formulated for their own populations have demonstrated that the latter yield more precise results.

This study confirms the effect of ethnicity on age estimation accuracy, as stepwise regression analysis showed that ethnicity and sex contributed significantly to the age estimation model. Therefore, ethnicity and sex were included in the age estimation equation, while tooth type was excluded. Additionally, a *t*-test showed a significant difference in PC/CVR values between Malays and Chinese population among 29-39 age group, which justified the inclusion of ethnicity in the age estimation equation.

The multiple linear regression analysis conducted in this study revealed a moderate correlation between age and the predictor variables for both the study and validation sample. Furthermore, the results demonstrated that the MAE values for the younger age groups were more favorable than those for the 50+ age group. This could be due to the lesser number of samples included in the current study amongst this particular age group [58]. Moreover, it must be noted that dentine deposition is slower in the elderly due to diminishing of blood, lymphatic, and nerve supply to the tooth [59]. In our study, MAE values ( $\leq 7.5$ ) showed higher accuracy than a previous study (8.41) [60]. At the same time, Anjani and Boedi [61], Kurniawan and Wibowo [62], and Song and Yang [63] reported higher accuracy than our study with  $MAE \leq 7.40$ .

In this study, a significant correlation was observed between the pulp chamber-to-crown volume ratio and chronological age in UM2s and LM2s among Malay and Chinese Malaysians. Even though research on age estimation using multi-rooted teeth remains scarce and is often time-intensive, different approaches could be

exceedingly advantageous, as previous studies have suggested that the accuracy of the estimated age is significantly improved by combining two or more age-determining techniques [64]. Despite these challenges, acquiring population-specific data across different ethnic groups is crucial for enhancing the accuracy of adult age estimation. This newly developed age estimation model holds potential for effective application among Malaysian mongoloids (Malays and Chinese) aged 18 to 50 years. The age estimate model for adults can help in solving criminal cases and other jurisdictions within Malaysia's legal justice system.

One limitation of this study is that the ethnicity of the samples was determined based on the patients' recorded names in the database. This limitation is a result of our study's retrospective design. Consequently, the rising trend of intermarriage among Malaysia's diverse ethnicities may have led to a biased sample distribution in a few cases. Also, in Malaysia, a person of Chinese or Indian descent who converts to Islam may adopt an Islamic/Arabic name. While determining ethnicity from names is a practical necessity in retrospective datasets lacking self-reported data, it introduces a known risk of non-differential misclassification bias. Within a multivariate regression framework, this can typically induce attenuation bias, which pushes the associated coefficients toward the null and underestimates true ethnic variance. However, the clinical and statistical impact of this attenuation is fundamentally limited by the shared phylogenetic and geographic ancestry of the studied populations. Because Malay and Chinese populations belong to the same broad East and Southeast Asian macro-ethnic lineage, the biological divergence in their dental development is relatively narrow.

Another concerning aspect is the inability of machine learning (ML) models to recognize different parts of teeth accurately; however, to reduce the effect of this error, the segmentation process in this study was a combination of automatic and manual segmentation. In addition, the 50+ group is poorly represented in this study sample due to increased loss of teeth in this age group.

## CONCLUSION

The primary objective of our research was to formulate a regression equation for age estimation within the Malaysian population by examining the correlation between chronological age and the PC/CVR in UM2 and LM2. The results revealed a moderate inverse correlation between chronological age and the PC/CVR. The findings indicated no statistically significant differences in the strength of the correlation between chronological age and the PC/CVR across sexes. This study substantiates that the PC/CVR serves as a reliable metric for estimating age in the Malaysian population aged 18 to 50 years.

## AUTHORS' CONTRIBUTIONS

The authors confirm their contribution to the paper as follows: R.I.A., P.A/L.K.N N., N.W.C.: Study conception and design; R.I.A., N.I.: Data collection; R.I.A., P.A/L.K.N

N., M.K.A.: Analysis and interpretation of results; R.I.A., N.W.C.: Draft manuscript. All authors reviewed the results and approved the final version of the manuscript.

## LIST OF ABBREVIATIONS

|                |                                                        |
|----------------|--------------------------------------------------------|
| CBCT           | = Cone Beam Computed Tomography                        |
| MIMICS         | = Materialise Interactive Medical Image Control System |
| 2D             | = Two-dimensional                                      |
| 3D             | = Three dimensional                                    |
| R <sup>2</sup> | = Coefficient of determination                         |
| R              | = Correlation coefficient                              |
| Micro- CT      | = Micro Computed Tomography                            |
| SPSS           | = Statistical package for the social sciences          |
| ICC            | = Intraclass correlation coefficient                   |
| MAE            | = Mean absolute error                                  |
| DICOM          | = Digital imaging and communications in medicine       |
| PTV            | = Pulp tooth volume                                    |
| PCV            | = pulp chamber volume                                  |
| PC/CVR         | = Pulp chamber/crown volume ratio                      |
| CEJ            | = cemento-enamel junction                              |
| UM2            | = maxillary second molar                               |
| LM2            | = mandibular second molar                              |

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The ethical approval for this retrospective study was obtained from the University of Malaya Medical Ethics Committee (Ref. No: DF OS2002/0003(L)).

## HUMAN AND ANIMAL RIGHTS

This study was conducted in accordance with ethical principles of the Declaration of Helsinki.

## CONSENT FOR PUBLICATIONS

Patients were well aware that the data from their CBCT scans could be utilized for research and education at the faculty, and consented to their use.

## STANDARDS OF REPORTING

STROBE guidelines were followed.

## AVAILABILITY OF DATA AND MATERIALS

The data and supportive information is available within the article.

## FUNDING

None.

## CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

## ACKNOWLEDGEMENTS

Declared none.

## REFERENCES

- [1] Willems G. A review of the most commonly used dental age estimation techniques. *J Forensic Odontostomatol* 2001; 19(1): 9-17. PMID: 11494678
- [2] Mani SA, Naing L, John J, Samsudin AR. Comparison of two methods of dental age estimation in 7-15-year-old Malays. *Int J Paediatr Dent* 2008; 18(5): 380-8. <http://dx.doi.org/10.1111/j.1365-263X.2007.00890.x> PMID: 18284472
- [3] Stavrianos C, Mastagas D, Stavrianou I, Karaïskou O. Dental age estimation of adults: A review of methods and principals. *Int J Res Med Sci* 2008; 2(5): 258-68.
- [4] Maled V, Manjunatha B, Patil K, Balaraj BM. The chronology of third molar root mineralization in south Indian population. *Med Sci Law* 2014; 54(1): 28-34. <http://dx.doi.org/10.1177/0025802413491557> PMID: 23966355
- [5] Pretty IA, Sweet D. A look at forensic dentistry - Part 1: The role of teeth in the determination of human identity. *Br Dent J* 2001; 190(7): 359-66. <http://dx.doi.org/10.1038/sj.bdj.4800972> PMID: 11338039
- [6] Alghali R, Kamaruddin AF, Mokhtar N. Dental age estimation: Comparison of reliability between Malay formula of Demirjian method and Malay formula of Cameriere method. *AIP Conf Proc* 2016; 1791(1): 020016. <http://dx.doi.org/10.1063/1.4968871>
- [7] Ubelaker DH, Khosrowshahi H. Estimation of age in forensic anthropology: Historical perspective and recent methodological advances. *Forensic Sci Res* 2019; 4(1): 1-9. <http://dx.doi.org/10.1080/20961790.2018.1549711> PMID: 30915413
- [8] Asif MK, Nambiar P, Mani SA, Ibrahim NB, Khan IM, Lokman NB. Dental age estimation in Malaysian adults based on volumetric analysis of pulp/tooth ratio using CBCT data. *Leg Med* 2019; 36: 50-8. <http://dx.doi.org/10.1016/j.legalmed.2018.10.005> PMID: 30415192
- [9] Gustafson G, Malmö DO. Age determination on teeth. *J Am Dent Assoc* 1950; 41(1): 45-54. <http://dx.doi.org/10.14219/jada.archive.1950.0132> PMID: 15428197
- [10] Ludlow JB, Timothy R, Walker C, Hunter R, Benavides E, Samuelson DB. Correction to effective dose of dental CBCT — A meta analysis of published data and additional data for nine CBCT units. *Dentomaxillofac Radiol* 2015; 44(7): 20159003. <http://dx.doi.org/10.1259/dmfr.20159003> PMID: 25874892
- [11] Pinchi V, Pradella F, Buti J, Baldinotti C, Focardi M, Norelli GA. A new age estimation procedure based on the 3D CBCT study of the pulp cavity and hard tissues of the teeth for forensic purposes: A pilot study. *J Forensic Leg Med* 2015; 36: 150-7. <http://dx.doi.org/10.1016/j.jflm.2015.09.015> PMID: 26458182
- [12] Yousefi F, Mohammadi Y, Ahmadvand M, Razaghi P. Dental age estimation using cone-beam computed tomography: A systematic review and meta-analysis. *Imaging Sci Dent* 2023; 53(2): 91-9. <http://dx.doi.org/10.5624/isd.20221226> PMID: 37405204
- [13] Merdjetio Boedi R, Shepherd S, Mânica S, Franco A. CBCT in dental age estimation: A systematic review and meta analysis. *Dentomaxillofac Radiol* 2022; 51(4): 20210335. <http://dx.doi.org/10.1259/dmfr.20210335> PMID: 34995103
- [14] Panchbhair AS. Dental radiographic indicators, a key to age estimation. *Dentomaxillofac Radiol* 2011; 40(4): 199-212. <http://dx.doi.org/10.1259/dmfr/19478385> PMID: 21493876
- [15] Cameriere R, Ferrante L, Cingolani M. Variations in pulp/tooth area ratio as an indicator of age: A preliminary study. *J Forensic Sci* 2004; 49(2): 1-3.

- <http://dx.doi.org/10.1520/JFS2003259> PMID: 15027553
- [16] Agematsu H, Sameda H, Hashimoto M, *et al.* Three-dimensional observation of decrease in pulp cavity volume using micro-CT: Age-related change. *Bull Tokyo Dent Coll* 2010; 51(1): 1-6. <http://dx.doi.org/10.2209/tdcpub.51.1> PMID: 20574128
  - [17] Vandevoort FM, Bergmans L, Cleynenbreugel JV, *et al.* Age calculation using X-ray microfocus computed tomographical scanning of teeth: A pilot study. *J Forensic Sci* 2004; 49(4): 1-4. <http://dx.doi.org/10.1520/JFS2004069> PMID: 15317195
  - [18] De Angelis D, Gaudio D, Guercini N, *et al.* Age estimation from canine volumes. *Radiol Med* 2015; 120(8): 731-6. <http://dx.doi.org/10.1007/s11547-015-0521-5> PMID: 25698302
  - [19] Biuki N, Razi T, Faramarzi M. Relationship between pulp-tooth volume ratios and chronological age in different anterior teeth on CBCT. *J Clin Exp Dent* 2017; 9(5): 0. <http://dx.doi.org/10.4317/jced.53654> PMID: 28512548
  - [20] Gulsahi A, Kulah CK, Bakirarar B, Gulen O, Kamburoglu K. Age estimation based on pulp/tooth volume ratio measured on cone-beam CT images. *Dentomaxillofac Radiol* 2018; 47(1): 20170239. <http://dx.doi.org/10.1259/dmfr.20170239> PMID: 28991500
  - [21] Yang F, Jacobs R, Willems G. Dental age estimation through volume matching of teeth imaged by cone-beam CT. *Forensic Sci Int* 2006; 159(Suppl. 1): S78-83. <http://dx.doi.org/10.1016/j.forsciint.2006.02.031> PMID: 16563678
  - [22] Star H, Thevissen P, Jacobs R, Fieuws S, Solheim T, Willems G. Human dental age estimation by calculation of pulp-tooth volume ratios yielded on clinically acquired cone beam computed tomography images of monoradicular teeth. *J Forensic Sci* 2011; 56(s1): S77-82. <http://dx.doi.org/10.1111/j.1556-4029.2010.01633.x> PMID: 21182523
  - [23] Asif MK, Nambiar P, Mani SA, Ibrahim NB, Khan IM, Sukumaran P. Dental age estimation employing CBCT scans enhanced with Mimics software: Comparison of two different approaches using pulp/tooth volumetric analysis. *J Forensic Leg Med* 2018; 54: 53-61. <http://dx.doi.org/10.1016/j.jflm.2017.12.010> PMID: 29324319
  - [24] Akbarizadeh F, Tabrizi BM, Fereidouni K, Mazidi MMK. Evaluation of the accuracy of age estimation based on ratio of pulp chamber to crown volume of the anterior teeth: A CBCT study. *Forensic Sci Int* 2025; 377: 112610. <http://dx.doi.org/10.1016/j.forsciint.2025.112610> PMID: 40857879
  - [25] De Micco F, Martino F, Velandia Palacio LA, Cingolani M, Campobasso CP. Third molar maturity index and legal age in different ethnic populations: Accuracy of Cameriere's method. *Med Sci Law* 2021; 61(1\_suppl): 105-12. <http://dx.doi.org/10.1177/0025802419900686> PMID: 33591873
  - [26] Jayaraman J, Wong HM, Roberts GJ, *et al.* Age estimation in three distinct east Asian population groups using southern Han Chinese dental reference dataset. *BMC Oral Health* 2019; 19(1): 242. <http://dx.doi.org/10.1186/s12903-019-0942-y> PMID: 31711475
  - [27] Talwar M, Chaudhry K, Choudhary A, Patnana AK, Vanga NRV, Lehl GK. A comparative evaluation of three different dental age estimation methods in India: A test of ethnic variability. *Int J Clin Pediatr Dent* 2020; 13(1): 16-20. <http://dx.doi.org/10.5005/jp-journals-10005-1708> PMID: 32581472
  - [28] World Medical Association. World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human participants. *JAMA* 2025; 333(1): 71-4. <http://dx.doi.org/10.1001/jama.2024.21972> PMID: 39425955
  - [29] Heidari S, Babor TF, De Castro P, Tort S, Curno M. Sex and Gender Equity in Research: Rationale for the SAGER guidelines and recommended use. *Res Integr Peer Rev* 2016; 1(1): 2. <http://dx.doi.org/10.1186/s41073-016-0007-6> PMID: 29451543
  - [30] Azim AA, Azim KA, Deutsch AS, Huang GTJ. Acquisition of anatomic parameters concerning molar pulp chamber landmarks using cone-beam computed tomography. *J Endod* 2014; 40(9): 1298-302. <http://dx.doi.org/10.1016/j.joen.2014.04.002> PMID: 25146010
  - [31] Alghali RI, Nambiar P, Ngeow WC, Ibrahim N, Asif MK, Al-Mouallad AS. Incidence and effect of an extra root on the crown morphometry of the maxillary second molars among Malaysian Mongoloids. *Folia Morphol* 2025; 84(3): 731-9. <http://dx.doi.org/10.5603/fm.101502> PMID: 39387396
  - [32] Sarment DP, Christensen AM. The use of cone beam computed tomography in forensic radiology. *J Forensic Radiol Imaging* 2014; 2(4): 173-81. <http://dx.doi.org/10.1016/j.jofri.2014.09.002>
  - [33] McKenna CJ, James H, Taylor JA, Townsend GC. Tooth development standards for South Australia. *Aust Dent J* 2002; 47(3): 223-7. <http://dx.doi.org/10.1111/j.1834-7819.2002.tb00333.x> PMID: 12405462
  - [34] Patel S. New dimensions in endodontic imaging: Part 2. Cone beam computed tomography. *Int Endod J* 2009; 42(6): 463-75. <http://dx.doi.org/10.1111/j.1365-2591.2008.01531.x> PMID: 19298576
  - [35] Di Lorenzo S, Zarin F, Pavone M, *et al.* Three-dimensional image guidance for diagnosis and treatment of adrenal disease: A systematic review. *Updates Surg* 2026; 78(1): 365-84. <http://dx.doi.org/10.1007/s13304-025-02386-9> PMID: 41068506
  - [36] Kvaal SI, Kolltveit KM, Thomsen IO, Solheim T. Age estimation of adults from dental radiographs. *Forensic Sci Int* 1995; 74(3): 175-85. [http://dx.doi.org/10.1016/0379-0738\(95\)01760-G](http://dx.doi.org/10.1016/0379-0738(95)01760-G) PMID: 7557754
  - [37] Olusegun IO, Gbolahan OO, Okoje VN, Sylvia M, Philip O. Comparative analyses of three radiographic dental age estimation methods amongst Nigerians. *Niger J Med* 2019; 28(4): 403-12. <http://dx.doi.org/10.4103/1115-2613.278632>
  - [38] Meinel A, Tangl S, Pernicka E, Fenes C, Watzek G. On the applicability of secondary dentin formation to radiological age estimation in young adults. *J Forensic Sci* 2007; 52(2): 438-41. <http://dx.doi.org/10.1111/j.1556-4029.2006.00377.x> PMID: 17316247
  - [39] Safaei A, Bagherpour A, Naseri S, Etemadi M, Khoshkhou H. Age estimation by evaluation of pulp chamber to crown volume of central incisor and first molar of maxilla, using Cone-Beam CT. *Heliyon* 2024; 10(21): e40022. <http://dx.doi.org/10.1016/j.heliyon.2024.e40022> PMID: 39568829
  - [40] Boyacioglu Erden H, Avcu N, Akcicek G, Akkaya N, Dural S. Volumetric pulp analysis of mandibular molars on CBCT as a supplementary indicator for age-related dental changes: A preliminary investigation in a Turkish population sample. *Diagnostics* 2026; 16(16): 2552. <http://dx.doi.org/10.3390/diagnostics16162552> PMID: 42650955
  - [41] Sindhuja N, Vardhan BGH, Sk G. Pulp Volume - An in-depth tool in age estimation- A comparative retrospective forensic based cone beam CT study. *J Indian Forensic Med* 2024; 46(2): 272-6. <http://dx.doi.org/10.48165/jiafm.2024.46.2.16>
  - [42] Vanessa T, Thais AC, Casimiro Abreu L, Luiz E, Eduardo E. Three-Dimensional Radiological Methods for Age Estimation in Adults by Using the Pulp/Tooth Relation: an Integrative Review. *Braz J Forensic Sci Med Law Bioeth* 2021; 10(2): 97-110. [http://dx.doi.org/10.17063/bjfs10\(2\)y202197-110](http://dx.doi.org/10.17063/bjfs10(2)y202197-110)
  - [43] Zhang Z, Yan C, Min Q, *et al.* Age estimation using pulp/enamel volume ratio of impacted mandibular third molars measured on CBCT images in a northern Chinese population. *Int J Legal Med* 2019; 133(6): 1925-33. <http://dx.doi.org/10.1007/s00414-019-02112-2> PMID: 31273446
  - [44] Porto LVMG, Celestino da Silva Neto J, Anjos Pontual A, Catunda RQ. Evaluation of volumetric changes of teeth in a Brazilian population by using cone beam computed tomography. *J Forensic Leg Med* 2015; 36: 4-9. <http://dx.doi.org/10.1016/j.jflm.2015.07.007> PMID: 26320003
  - [45] Tardivo D, Sastre J, Ruquet M, *et al.* Three-dimensional modeling of the various volumes of canines to determine age and sex: A preliminary study. *J Forensic Sci* 2011; 56(3): 766-70. <http://dx.doi.org/10.1111/j.1556-4029.2011.01720.x> PMID: 21361946
  - [46] Sue M, Oda T, Sasaki Y, Ogura I. Age-related changes in the pulp

- chamber of maxillary and mandibular molars on cone-beam computed tomography images. *Oral Radiol* 2018; 34(3): 219-23. <http://dx.doi.org/10.1007/s11282-017-0300-1> PMID: 30484030
- [47] Uğur Aydın Z, Bayrak S. Relationship between pulp tooth area ratio and chronological age using cone-beam computed tomography images. *J Forensic Sci* 2019; 64(4): 1096-9. <http://dx.doi.org/10.1111/1556-4029.13986> PMID: 30562410
- [48] Salemi F, Farhadian M, Askari Sabzkouhi B, Saati S, Nafisi N. Age estimation by pulp to tooth area ratio in canine teeth using cone-beam computed tomography. *Egypt J Forensic Sci* 2020; 10(1): 2. <http://dx.doi.org/10.1186/s41935-019-0176-9>
- [49] Aboshi H, Takahashi T, Komuro T. Age estimation using microfocus X-ray computed tomography of lower premolars. *Forensic Sci Int* 2010; 200(1-3): 35-40. <http://dx.doi.org/10.1016/j.forsciint.2010.03.024> PMID: 20381983
- [50] Yang Z, Fan L, Kwon K, et al. Age estimation for children and young adults by volumetric analysis of upper anterior teeth using cone-beam computed tomography data. *Folia Morphol* 2020; 79(4): 851-9. <http://dx.doi.org/10.5603/FM.a2020.0004> PMID: 31930465
- [51] Türk E, Görmez Ö, Yıldırım D. Age estimation of adults from cone beam computed tomography images. *Eur Ann Dent Sci* 2022; 49(Suppl 1): 37-41. <http://dx.doi.org/10.52037/eads.2022.0020>
- [52] Zhan M, Chen X, Shi L, et al. Age estimation in Western Chinese adults by pulp-tooth volume ratios using cone-beam computed tomography. *Aust J Forensic Sci* 2021; 53(6): 681-92. <http://dx.doi.org/10.1080/00450618.2020.1729415>
- [53] Krishnapriya BDSJ, Debta FM, Agarwal K, Debta P, Bohidar P, Mishra E. Assessment of dental age using CBCT-based pulp/tooth volume ratio of mandibular premolars in Odisha population for forensic age estimation - An *in vitro* study. *J Indian Acad Oral Med Radiol* 2023; 35(4): 562-6. [http://dx.doi.org/10.4103/jiaomr.jiaomr\\_6\\_23](http://dx.doi.org/10.4103/jiaomr.jiaomr_6_23)
- [54] Prakash P. Age estimation in Indian soldiers: Maxillary anterior tooth pulp/tooth volume ratio analysis with cone beam computed tomography. *Med J Armed Forces India* 2024; 80(4): 458-65. <http://dx.doi.org/10.1016/j.mjafi.2024.04.016> PMID: 39071747
- [55] Elgazzar FM, Elboraey MO, El-Sarnagawy GN. The accuracy of age estimation from pulp chamber/crown volume ratio of canines obtained by cone beam computed tomography images: An Egyptian study. *Egypt J Forensic Sci* 2020; 10(1): 40. <http://dx.doi.org/10.1186/s41935-020-00212-4>
- [56] Hatice BD, Nihal A, Nursel A, Humeyra Ozge Y, Goksuluk D. Applicability of Cameriere's and Drusini's age estimation methods to a sample of Turkish adults. *Dentomaxillofac Radiol* 2017; 46(7): 20170026. <http://dx.doi.org/10.1259/dmfr.20170026> PMID: 28707524
- [57] Molina A, Bravo M, Fonseca GM, Márquez-Grant N, Martín-de-las-Heras S. Dental age estimation based on pulp chamber/crown volume ratio measured on CBCT images in a Spanish population. *Int J Legal Med* 2021; 135(1): 359-64. <http://dx.doi.org/10.1007/s00414-020-02377-y> PMID: 32676887
- [58] Vossoughi M, Movahhedian N, Ghafoori A. The impact of age mimicry bias on the accuracy of methods for age estimation based on Kvaal's pulp/tooth ratios: A bootstrap study. *Int J Legal Med* 2022; 136(1): 269-78. <http://dx.doi.org/10.1007/s00414-021-02651-7> PMID: 34291317
- [59] Morse DR. Age-related changes of the dental pulp complex and their relationship to systemic aging. *Oral Surg Oral Med Oral Pathol* 1991; 72(6): 721-45. [http://dx.doi.org/10.1016/0030-4220\(91\)90019-9](http://dx.doi.org/10.1016/0030-4220(91)90019-9) PMID: 1812456
- [60] Yüksel İB, Altındağ A. Age estimation based on pulp/tooth volume ratio using CBCT: A study on mandibular canines. *BMC Oral Health* 2025; 25(1): 1248. <http://dx.doi.org/10.1186/s12903-025-06659-y> PMID: 40713693
- [61] Anjani KG, Merdietio Boedi R, Sam B, Oscandar F. Dental age estimation in Indonesian adults: An investigation of the maxillary canine pulp-to-tooth volume ratio using cone-beam computed tomography. *Imaging Sci Dent* 2023; 53(3): 221-8. <http://dx.doi.org/10.5624/isd.20230104> PMID: 37799738
- [62] Kurniawan A, Wibowo MRR, Chusida A, et al. Adults' dental age estimation by Cameriere's method using mandibular canines' pulp/tooth ratio in Surabaya, Indonesia. *Bull Int Assoc Paleodont* 2022; 16(1): 8-16.
- [63] Song Y, Yang H, Ge Z, Du H, Li G. Age estimation based on 3D pulp segmentation of first molars from CBCT images using U-Net. *Dentomaxillofac Radiol* 2023; 52(7): 20230177. <http://dx.doi.org/10.1259/dmfr.20230177> PMID: 37427595
- [64] Nayyar A, Babu BA, Krishnaveni B, Devi MV, Gayitri HC. Age estimation: Current state and research challenges. *Yixue Yanjiu Zazhi* 2016; 36(6): 209-16. <http://dx.doi.org/10.4103/1011-4564.196348>

**DISCLAIMER:** Please note that this article is currently in the 'Early view' stage and is not the final 'Version of Record'. While it has been accepted, copy-edited, and formatted, however, it is still undergoing proofreading and corrections by the authors. Therefore, the text may still change before the final publication. Although 'Early view' may not have all bibliographic details available, the DOI and the year of online publication can still be used to cite them. The article title, DOI, publication year, and author(s) should all be included in the citation format. Once the 'Final Version of Record' becomes available the 'Early view' will be replaced by that.