The Open Dentistry Journal
DOI: 10.2174/0118742106294926101609275O, 2026, 20, e187421062949
ISSN: 1874-2106
1

RESEARCH ARTICLE **OPEN ACCESS**

Clinical and Radiographic Outcomes of Pulpectomy in Primary Second Molars with a History of Abscess or Facial Cellulitis: A Comparative Clinical Study

Ziya Ebrahim Adhami[1], Mahsa Karkhaneh[1,*], Ali Vafaei[1], Naimeh Teimoory[1] and Mehdi Dehghan[2]

[1]*Department of Pediatric Dentistry, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran*
[2]*Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran*

Abstract:

Introduction: Odontogenic infections originating from primary teeth remain a frequent cause of pain, facial swelling, and systemic complications in children. Although pulpectomy is the treatment of choice for necrotic primary teeth, the influence of a previous history of abscess or facial cellulitis on treatment prognosis remains controversial. This study aimed to compare the clinical and radiographic success of pulpectomy in primary second molars with a history of abscess or facial cellulitis and those with irreversible pulpitis without prior infection.

Materials and Methods: This clinical study included 58 primary second molars in children aged 3–9 years (n=58). Teeth were allocated into a case group (necrotic teeth with a history of acute or chronic abscess or facial cellulitis, treated with two-visit pulpectomy) and a control group (vital teeth with irreversible pulpitis, treated with single-visit pulpectomy). All canals were instrumented manually, obturated with zinc oxide–eugenol, and restored with stainless steel crowns. Clinical and radiographic outcomes were evaluated after 6 months. Fisher's exact test was used for statistical analysis. Fisher's exact test was used for statistical analysis, and Relative Risk (RR) and Absolute Risk Reduction (ARR) were calculated. A *p*-value less than 0.05 was considered significant.

Results: Radiographic success was 89.7% in the case group and 88.9% in the control group (p = 1.00). Clinical success was 96.6% and 96.3%, respectively (p = 1.00). No statistically significant differences were observed between groups in any radiographic or clinical subgroup outcomes.

Discussion: Pulpectomy in necrotic primary second molars with a history of abscess or facial cellulitis demonstrates clinical and radiographic outcomes comparable to those of vital teeth.

Conclusion: When infection is properly managed and standard endodontic and restorative protocols are followed, prior odontogenic infection does not negatively affect prognosis.

Keywords: Pulpectomy, Primary teeth, Dental abscess, Facial cellulitis, Pediatric dentistry.

© 2026 The Author(s). Published by Bentham Open.
This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: https://creativecommons.org/licenses/by/4.0/legalcode. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Department of Pediatric Dentistry, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran; Email: mahsakarkhane2377@gmail.com

Cite as: Adhami Z, Karkhaneh M, Vafaei A, Teimoory N, Dehghan M. Clinical and Radiographic Outcomes of Pulpectomy in Primary Second Molars with a History of Abscess or Facial Cellulitis: A Comparative Clinical Study. Open Dent J, 2026; 20: e187421062949. http://dx.doi.org/10.2174/0118742106294926101609275O

Received: February 02, 2026
Revised: August 19, 2026
Accepted: September 03, 2026
Published: November 10, 2026

Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Dental caries remains one of the most prevalent chronic diseases in childhood and frequently progresses to pulpal involvement due to the structural characteristics of primary teeth. Untreated pulpal disease may result in pain, impaired mastication, sleep disturbances, and reduced quality of life [1]. In advanced stages, odontogenic infections such as dentoalveolar abscesses and facial cellulitis may develop, which can progress rapidly in children and be associated with serious systemic complications [2].

Primary teeth play a critical role in maintaining arch length, guiding eruption of permanent successors, preserving oral function, and supporting speech and facial esthetics. Premature loss of primary molars, particularly second molars, may lead to malocclusion and functional problems [3]. Therefore, preserving primary teeth until their natural exfoliation is a fundamental principle in pediatric dentistry [4].

Pulpectomy is recommended by the American Academy of Pediatric Dentistry (AAPD) as the treatment of choice for necrotic primary teeth when adequate root structure remains [5, 6]. Despite this recommendation, clinicians often hesitate to perform pulpectomy in teeth with a history of abscess or facial cellulitis due to concerns about persistent infection, compromised healing, and potential effects on the permanent successor [7].

In a study by Mendoza *et al*., pulpectomized teeth with necrosis were studied, and complete resolution of pain, swelling, and fistula was observed within 6 months, as well as progressive healing of the radiolucent lesion on radiographs [8]. In a study by Shenkin, 308 pulpectomized teeth with irreversible pulpitis were treated with zinc oxide eugenol root filling material, and 100% clinical success and 93.5% radiological success were observed [9]. In a study by Gupta *et al*., 42 necrotic teeth were randomly assigned to two groups, and two root filling materials, zinc oxide eugenol and Metapex, were compared, and the results were almost similar for them [10].

According to AAPD, untreated dental caries and infections affect the ability to eat, resulting in poor nutritional intake and stunted or reduced growth in children, especially in low-income families [9]. Treatment of dental caries and infections increases growth rates and improves the quality of life of millions of children [11]. Odontogenic infections differ in children and adults. The course of facial infection in children is dynamic and more acute, associated with faster spread to deeper spaces, and leads to more systemic symptoms such as fever, dehydration, and impaired respiratory function in children; therefore, prompt treatment is necessary [12].

On the other hand, premature tooth loss due to necrosis and dental infection leads to malocclusion and functional, aesthetic, and phonetic problems, which can be prevented by preserving the tooth with timely pulpectomy [13]. Therefore, the primary objective of this study was to compare the clinical and radiographic success rates of pulpectomy in primary second molars with a history of dentoalveolar abscess or facial cellulitis (treated with a two-visit protocol) against vital teeth with irreversible pulpitis (treated with a single-visit protocol) over a 6-month follow-up period. The secondary objective was to evaluate specific failure criteria, including periapical/furcation radiolucencies, pathological resorption, recurrent pain, mobility, and sinus tract formation. We hypothesized that necrotic primary second molars with a prior history of acute or chronic abscess or facial cellulitis would demonstrate clinical and radiographic success rates comparable to those of vital teeth when proper disinfection and coronal sealing protocols are applied (null hypothesis).

2. MATERIALS AND METHODS

2.1. Sample Size

The sample size was calculated based on expecting a clinical/radiographic success rate of approximately 90% in primary molar pulpectomies according to previous literature. To detect a clinically relevant difference of 25% in success rates between the two groups with a statistical power of 80% ($\beta=0.20$) and a two-tailed significance level of 5% ($\alpha=0.05$), a minimum of 26 teeth per group was required. Anticipating a potential drop-out rate of approximately 10% during follow-up, 29 teeth were initially enrolled in each group (total N=58).

2.2. The Population and the Study Design

A total of 58 children aged 3–9 years requiring pulpectomy of a primary second molar were recruited. A total of 58 teeth were included and allocated into two groups. The case group consisted of 29 necrotic primary second molars with a documented history of acute or chronic abscess or facial cellulitis. The control group consisted of 29 vital primary second molars diagnosed with irreversible pulpitis.

2.3. Inclusion Criteria

Inclusion criteria were restorable teeth with more than two-thirds of root length remaining and absence of extensive internal or external root resorption. Teeth with lesions extending to the permanent tooth germ, advanced physiologic resorption, pathological mobility, or previous pulp therapy were excluded.

2.4. Procedures

All procedures were performed under local anesthesia and rubber dam isolation. Access cavities were prepared, and working length was determined radiographically. Canals were instrumented manually using K-files up to size 30 with copious saline irrigation. In the case group, treatment was completed in two visits, and children presenting with facial cellulitis received systemic antibiotics prior to endodontic treatment. In the control group, pulpectomy was completed in a single visit using the same protocol. In both groups, canals were obturated with zinc oxide–eugenol paste and restored with stainless steel crowns cemented with glass ionomer cement.

Given the anatomical complexity of primary molar root canal systems, including accessory canals and physiological resorption patterns, particular attention was paid to conservative instrumentation to prevent over-preparation and extrusion of obturation material beyond the apex. This approach was especially critical in teeth with previous infection, where periapical tissues may already be compromised. The two-visit protocol selected for teeth with a history of abscess or facial cellulitis was designed to allow additional microbial reduction and periapical tissue recovery prior to obturation. The use of an interim intracanal medicament and delayed obturation was intended to enhance bacterial suppression and reduce the risk of persistent infection. Antibiotic Therapy (AT) in cases presenting with facial cellulitis was prescribed strictly according to clinical indications and current pediatric guidelines, emphasizing that systemic antibiotics were used as an adjunct to, rather than a substitute for, definitive endodontic treatment.

To enhance methodological consistency and minimize operator-related bias, all clinical procedures in both study groups were performed according to a standardized and uniform protocol. The same instrumentation sequence, irrigation regimen, obturation material (zinc oxide–eugenol), and restorative approach were applied to all treated teeth to control for potential confounding variables. Manual instrumentation was selected to ensure precise canal debridement in primary teeth with complex root morphology. Repeated irrigation with normal saline and sodium hypochlorite was employed to reduce microbial load and remove necrotic debris, particularly in teeth with a previous history of infection.

All treated teeth were definitively restored with stainless steel crowns to ensure optimal coronal sealing and to minimize microleakage, which has been identified as a critical determinant of pulpectomy success in primary teeth. This standardized coronal restoration protocol was implemented to eliminate restoration-related variability between groups. Clinical and radiographic evaluations were independently performed by two experienced pediatric dentists who were blinded to the pulp status and group allocation of the teeth. This blinding process was designed to reduce assessment bias and improve the internal validity of the study.

2.5. Statistical Analysis

Clinical and radiographic evaluations were performed after 6 months by two independent pediatric dentists blinded to group allocation. Fisher's exact test was used for statistical analysis, and Relative Risk (RR) and Absolute Risk Reduction (ARR) were calculated. A *p*-value less than 0.05 was considered significant.

3. RESULTS

At the 6-month follow-up, complete data were available for 56 teeth (27 control, 29 case). The median age of participants was 5 years (interquartile range: 5–6), with a mean age of 5.59 ± 1.14 years. Sex distribution was identical between groups (Table **1**).

Table 1. Demographic characteristics.

Variable	Total (n=58)	Control (n=29)	Case (n=29)
Age, mean ± SD	5.59 ± 1.14	5.79 ± 1.01	5.38 ± 1.24
Age, median (IQR)	5 (5–6)	6 (5–6.5)	5 (5–6)
Male, n (%)	32 (55.2)	16 (55.2)	16 (55.2)
Female, n (%)	26 (44.8)	13 (44.8)	13 (44.8)

Radiographic success was observed in 26 of 29 teeth (89.7%) in the case group and 24 of 27 teeth (88.9%) in the control group, with no statistically significant difference (p = 1.00; RR = 1.008). Table **2** shows the overall radiographic success.

Table 2. Overall radiographic success.

Outcome	Control (n=27)	Case (n=29)	*p*-value
Success	24 (88.9%)	26 (89.7%)	1.00
Failure	3 (11.1%)	3 (10.3%)	—

Clinical success was achieved in 28 of 29 teeth (96.6%) in the case group and 26 of 27 teeth (96.3%) in the control group (p = 1.00; RR = 1.002). Table **3** shows the overall clinical success.

Table 3. Overall clinical success.

Outcome	Control (n=27)	Case (n=29)	*p*-value
Success	26 (96.3%)	28 (96.6%)	1.00
Failure	1 (3.7%)	1 (3.4%)	—

Osseous radiolucency improvement was observed in 41.4% of the case group and 18.5% of the control group (p = 0.08). Periodontal ligament widening was detected in 10.3% of the case group and 11.1% of the control group (p = 1.00). Crypt involvement was observed in 6.9% of the case group and 3.7% of the control group (p = 1.00). Root resorption occurred in 10.3% of the case group and 11.1% of the control group (p = 1.00). Table **4** shows radiographic subgroup outcomes.

Clinically, pain and mobility were each observed in one tooth (3.7%) in the control group and none in the case group (p = 0.48). Gingival enlargement was detected in one tooth (3.4%) in the case group (p = 0.48). Erythema was present in one tooth in each group (p = 1.00). No teeth in either group exhibited percussion sensitivity (Table **5**).

Table 4. Radiographic subgroup outcomes.

Parameter	Control n (%)	Case n (%)	*p*-value
Osseous radiolucency	5 (18.5)	12 (41.4)	0.08
PDL widening	3 (11.1)	3 (10.3)	1.00
Crypt involvement	1 (3.7)	2 (6.9)	1.00
Root resorption	3 (11.1)	3 (10.3)	1.00

Table 5. Clinical subgroup outcomes.

Parameter	Control n (%)	Case n (%)	*p*-value
Pain	1 (3.7)	0 (0)	0.48
Mobility	1 (3.7)	0 (0)	0.48
Percussion sensitivity	0 (0)	0 (0)	—
Gingival enlargement	0 (0)	1 (3.4)	0.48
Erythema	1 (3.7)	1 (3.4)	1.00

4. DISCUSSION

The primary aim of the present study was to evaluate whether a previous history of odontogenic infection, including dental abscess or facial cellulitis, influences the clinical and radiographic outcomes of pulpectomy in primary second molars. The results demonstrated that pulpectomy performed in necrotic primary teeth with prior infection achieved success rates comparable to those of vital teeth diagnosed with irreversible pulpitis. The present study demonstrated that pulpectomy in necrotic primary second molars with a history of abscess or facial cellulitis resulted in clinical and radiographic success rates comparable to those of vital primary second molars with irreversible pulpitis. Both groups exhibited high success rates, and no statistically significant differences were identified in overall outcomes or in any evaluated clinical or radiographic subgroups.

The radiographic success rates observed in this study are consistent with those reported in previous systematic reviews, which describe success rates ranging from 70% to 94% for pulpectomy in primary teeth [14]. Importantly, the similarity between groups suggests that preoperative infection status does not inherently compromise radiographic healing when proper canal debridement and coronal sealing are achieved. This finding is in agreement with studies by Holland *et al*. [15] and Khan *et al*., who reported that the presence of necrosis or previous infection did not adversely affect pulpectomy outcomes [16].

In contrast, some studies have reported lower success rates in infected primary teeth. Differences between those findings and the present results may be explained by variations in treatment protocols, particularly irrigation methods [17], obturation materials, and the quality of coronal restorations [17]. In the present study, all teeth were restored with stainless steel crowns, which likely minimized microleakage and contributed to the favorable outcomes [17]. Some studies have also shown that children have the highest rates of bone repair [18, 19].

Although osseous radiolucency improvement was numerically higher in the case group, this difference was not statistically significant. From a biological perspective, removal of necrotic tissue and microbial load may activate reparative pathways and osteoblastic activity, leading to radiographic evidence of healing [20, 21]. The comparable radiographic healing patterns observed in both groups may be explained by the high regenerative capacity of pediatric periapical tissues. Children exhibit accelerated bone turnover and enhanced inflammatory resolution, which may compensate for the initial severity of infection once the source of microbial irritation is eliminated [5, 22, 23].

In a study, Abanto *et al*. reported that after pulpectomy, clinical symptoms resolved in more than 94% of children within the first months [5]. Samir *et al*. also stated that the type of pulp diagnosis did not significantly affect the appearance of pain [24]. This result is consistent with our data.

However, there are studies with inconsistent results. A study by Mendoza *et al*. on pulpectomy treatment of necrotic primary teeth with a history of abscesses showed that treatment of these teeth is less successful than for healthy teeth. In this study, the success rate of treatment in primary teeth with a history of abscess was reported to be only 70%, while this rate was 90% in healthy teeth [8]. In a study by Shenkin *et al*., it was shown that primary teeth with a history of abscess or facial cellulitis are generally more difficult to treat and their success rate is significantly lower than in healthy teeth or teeth with a viable pulp. This study also concluded that additional treatments and continuous follow-up are necessary for primary teeth with a history of infection [9]. A study by Gupta *et al*. conducted on primary teeth with irreversible pulpitis and necrotic primary teeth states that pulpectomy treatment is less successful in primary teeth with a history of infection than in healthy teeth. Also, in necrotic teeth, the rate of infection recurrence and the need for follow-up treatments were observed to be higher [10]. In the study of Bawazir *et al*., they investigated the success of pulpectomy treatment in necrotic primary teeth using zinc oxide eugenol and showed that the success rate of treatment in primary teeth with a history of infection was significantly reduced compared to healthy teeth [25].

These findings of the current study support the hypothesis that when adequate infection control, thorough canal debridement, and appropriate coronal restoration are achieved, a history of odontogenic infection does not adversely affect pulpectomy prognosis. This is particularly relevant in pediatric patients, where bone healing capacity and regenerative potential are generally higher than in adults. The absence of statistically significant differences in radiographic subgroups, including osseous radiolucency, periodontal ligament widening, crypt involvement, and root resorption, suggests that periapical healing is more closely associated with treatment quality rather than the initial vitality status of the pulp [26]. Proper elimination of necrotic tissue and microbial contamination appears to promote favorable periapical repair even in previously infected teeth [27, 28]. From a clinical perspective, the high success rates observed in both groups indicate effective resolution of inflammatory signs such as pain, mobility, gingival changes, and percussion sensitivity. These outcomes reinforce the role of pulpectomy as a reliable and conservative treatment option for maintaining infected primary molars until their natural exfoliation.

Variability among previously published studies may also be attributed to differences in follow-up duration, operator experience, obturation techniques, and coronal restoration quality [29, 30]. In the present study, the uniform use of stainless steel crowns likely contributed to the high overall success rates by preventing coronal leakage and reinfection.

5. LIMITATIONS

Limitations of this study include the 6-month follow-up period, absence of Cone Beam Computed Tomography (CBCT) imaging, and small sample sizes in some subgroups. Longer follow-up and advanced imaging may provide additional insight into long-term outcomes.

CONCLUSION

Based on the findings of this study, a previous history of dental abscess or facial cellulitis should not be considered a contraindication for pulpectomy in primary second molars. When infection is appropriately managed and standard endodontic and restorative principles are strictly followed, favorable clinical and radiographic outcomes can be achieved. The results of this study support a more conservative treatment approach and may assist pediatric dentists in avoiding unnecessary extraction of infected primary molars, thereby preventing subsequent orthodontic, functional, and psychosocial complications associated with premature tooth loss. From a broader clinical perspective, preserving infected primary molars may reduce the need for space maintainers, decrease the risk of malocclusion, and support normal oral development, reinforcing the importance of conservative pulp therapy in pediatric dentistry.

AUTHORS' CONTRIBUTIONS

The authors confirm their contributions to the paper as follows: Z.E.A., M.K.: Conceptualization; A.V., N.T., M.D.: Data curation; M.K.: Formal analysis; Z.E.A.: Investigation; Z.E.A.: Methodology; Z.E.A.: Project administration; M.K.: Resources; Z.E.A.: Supervision; A.V., Z.E.A.: Validation; M.K., M.D.: Writing-original draft; Z.E.A., A.V.: Writing-review & editing.

LIST OF ABBREVIATIONS

AAPD = American Academy of Pediatric Dentistry

AT = Antibiotic therapy

CBCT = Cone Beam Computed Tomography

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This clinical study was conducted at the Faculty of Dentistry, Tabriz University of Medical Sciences, following approval by the institutional ethics committee (IR.TBZMED.REC.1404.538).

HUMAN AND ANIMAL RIGHTS

All procedures involving human participants were conducted in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

Written informed consent was obtained from parents or legal guardians.

STANDARDS OF REPORTING

STROBE guidelines were followed.

AVAILABILITY OF DATA AND MATERIALS

The data and supporting information is available within the article.

FUNDING

This paper was granted by the Vice Chancellor for Research. Tabriz University of Medical Sciences (grant number: 75996).

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

This paper was written based on a thesis granted by the Vice Chancellor for Research. Tabriz University of Medical Sciences (grant number: 75996), which is acknowledged.

REFERENCES

[1] Seow WK. Diagnosis and management of unusual dental abscesses in children. Aust Dent J 2003; 48(3): 156-68. http://dx.doi.org/10.1111/j.1834-7819.2003.tb00026.x PMID: 14640368

[2] Mathur VP, Dhillon JK. Dental caries: A disease which needs attention. Indian J Pediatr 2018; 85(3): 202-6. http://dx.doi.org/10.1007/s12098-017-2381-6 PMID: 28643162

[3] Hanafi L. An approach of preserving a mandibular primary second molar by a hemisection procedure: A case report with 36-month follow up. Heliyon 2022; 8(9): e10655. http://dx.doi.org/10.1016/j.heliyon.2022.e10655 PMID: 36158099

[4] Moskovitz M, Tickotsky N. Pulpectomy and Root Canal Treatment (RCT) in primary teeth: Techniques and materials. In: Fuks A, Peretz B, Eds. Pediatric Endodontics. Cham: Springer 2016; pp. 71-101. http://dx.doi.org/10.1007/978-3-319-27553-6_6

[5] Abanto J, Tsakos G, Olegário IC, *et al*. Impact of pulpectomy *versus* tooth extraction in children's oral health-related quality of life: A randomized clinical trial. Community Dent Oral Epidemiol 2024; 52(1): 13-23. http://dx.doi.org/10.1111/cdoe.12895 PMID: 37519111

[6] Padmawar N, Pawar N, Tripathi V, Banerjee S, Tyagi G, Joshi SR. Comparative analysis of rotary *versus* manual instrumentation in paediatric pulpectomy procedures: A systematic review and meta-analysis. Aust Endod J 2025; 51(1): 181-96. http://dx.doi.org/10.1111/aej.12899 PMID: 39494967

[7] Zhang Y, Sheng M. Clinical and radiographic evaluation of regenerative endodontic procedures (REPs) with or without concentrated growth factor (CGF) as scaffolds for non-vital immature mandibular premolars. J Clin Pediatr Dent 2024; 48(4): 168-75. http://dx.doi.org/10.22514/jocpd.2024.090 PMID: 39087227

[8] Mendoza AM, Reina JE, Garcia-Godoy F. Evolution and prognosis of necrotic primary teeth after pulpectomy. Am J Dent 2010;

23(5): 265-8.
PMID: 21207793
[9] Shenkin JD. Pulpectomy of necrotic primary teeth may be an effective tool in managing the primary dentition. J Evid Based Dent Pract 2012; 12(1): 39-40.
http://dx.doi.org/10.1016/j.jebdp.2011.12.014 PMID: 22326161
[10] Gupta S, Das G. Clinical and radiographic evaluation of zinc oxide eugenol and metapex in root canal treatment of primary teeth. J Indian Soc Pedod Prev Dent 2011; 29(3): 222-8.
http://dx.doi.org/10.4103/0970-4388.85829 PMID: 21985878
[11] Sheiham A. Dental caries affects body weight, growth and quality of life in pre-school children. Br Dent J 2006; 201(10): 625-6.
http://dx.doi.org/10.1038/sj.bdj.4814259 PMID: 17128231
[12] Słotwińska-Pawlaczyk A, Orzechowska-Wylęgała B, Latusek K, Roszkowska AM. Analysis of the clinical status and treatment of facial cellulitis of odontogenic origin in pediatric patients. Int J Environ Res Public Health 2023; 20(6): 4874.
http://dx.doi.org/10.3390/ijerph20064874 PMID: 36981782
[13] Tiwari S, Kulkarni P, Agrawal N, Kumar A, Umarekar P, Bhargava S. Clinical outcome of direct pulp therapy in primary teeth: A systematic review and meta-analysis. J Indian Soc Pedod Prev Dent 2022; 40(2): 105-11.
http://dx.doi.org/10.4103/jisppd.jisppd_210_22 PMID: 35859400
[14] Ahmed HMA. Pulpectomy procedures in primary molar teeth. Eur J Gen Dent 2014; 3(1): 3-10.
http://dx.doi.org/10.4103/2278-9626.126201
[15] Holland M, Carrico C, Williams T. Pediatric dentists' treatment planning decisions for class II carious lesions in primary molars. ASDC J Dent Child 2018; 85(1): 16-22.
PMID: 29663970
[16] Khan MK, Sharma DS, Jindal MK. Laser photobiomodulation as noninvasive therapeutic modality in pediatric endodontics: A comprehensive review. J Sci Soc 2024; 51(2): 147-52.
http://dx.doi.org/10.4103/jss.jss_48_23
[17] Latham J, Fong H, Jewett A, Johnson JD, Paranjpe A. Disinfection efficacy of current regenerative endodontic protocols in simulated necrotic immature permanent teeth. J Endod 2016; 42(8): 1218-25.
http://dx.doi.org/10.1016/j.joen.2016.05.004 PMID: 27372159
[18] Clark EM, Tobias JH, Ness AR. Association between bone density and fractures in children: A systematic review and meta-analysis. Pediatrics 2006; 117(2): e291-7.
http://dx.doi.org/10.1542/peds.2005-1404 PMID: 16452336
[19] Lindaman LM. Bone healing in children. Clin Podiatr Med Surg 2001; 18(1): 97-108.
http://dx.doi.org/10.1016/S0891-8422(23)01169-2 PMID: 11344982
[20] Segura-Sampedro JJ, Jiménez-Giménez C, Jane-Salas E, *et al*. Periapical and endodontic status of patients with inflammatory bowel disease: Age- and sex-matched case–control study. Int Endod J 2022; 55(7): 748-57.
http://dx.doi.org/10.1111/iej.13747 PMID: 35403728
[21] Piras V, Usai P, Mezzena S, *et al*. Prevalence of apical periodontitis in patients with inflammatory bowel diseases: A retrospective clinical study. J Endod 2017; 43(3): 389-94.
http://dx.doi.org/10.1016/j.joen.2016.11.004 PMID: 28231978
[22] Mathews M, Boykins J, Schoenthaler AM, Gold HT, Burns LE. Access to care for the endodontic treatment of pediatric patients: A qualitative study. Int J Paediatr Dent 2026; 36(2): 311-22.
http://dx.doi.org/10.1111/ipd.70054 PMID: 41225291
[23] Chen SY, Wang HL, Glickman GN. The influence of endodontic treatment upon periodontal wound healing. J Clin Periodontol 1997; 24(7): 449-56.
http://dx.doi.org/10.1111/j.1600-051X.1997.tb00211.x PMID: 9226384
[24] Samir PV, Mahapatra N, Dutta B, Bagchi A, Dhull KS, Verma RK. A Correlation between clinical classification of dental pulp and periapical diseases with its patho physiology and pain pathway. Int J Clin Pediatr Dent 2023; 16(4): 639-44.
http://dx.doi.org/10.5005/jp-journals-10005-2636 PMID: 37731799
[25] Bawazir OA, Salama FS. Clinical evaluation of root canal obturation methods in primary teeth. Pediatr Dent 2006; 28(1): 39-47.
PMID: 16615374
[26] Zhang Y, Yang J, Cui EH, *et al*. Impact on pulp vitality of affected teeth after decompression for jaw cystic lesions. J Endod 2025; 51(9): 1170-6.
http://dx.doi.org/10.1016/j.joen.2025.05.021 PMID: 40451423
[27] Duncan HF. Present status and future directions - Vital pulp treatment and pulp preservation strategies. Int Endod J 2022; 55(Suppl 3): 497-511.
http://dx.doi.org/10.1111/iej.13688 PMID: 35080024
[28] Asgary S, Nosrat A. Vital Pulp Therapy: Evidence-based techniques and outcomes. Iran Endod J 2025; 20(1): e2.
http://dx.doi.org/10.22037/iej.v20i1.47141 PMID: 39935864
[29] Thakur B, Bhardwaj A, Wahjuningrum DA, *et al*. Incidence of post-operative pain following a single-visit pulpectomy in primary molars employing adaptive, rotary, and manual instrumentation: A randomized clinical trial. Medicina 2023; 59(2): 355.
http://dx.doi.org/10.3390/medicina59020355 PMID: 36837556
[30] Shah HS, Patil VM, Kamath AP, Mathur AA. Comparative evaluation of instrumentation time, obturation time, and radiographic quality of obturation using two rotary systems and manual technique for primary molar pulpectomies – *In vivo* study. Contemp Clin Dent 2021; 12(1): 55-62.
http://dx.doi.org/10.4103/ccd.ccd_83_20 PMID: 33967539

DISCLAIMER: Please note that this article is currently in the 'Early view' stage and is not the final 'Version of Record'. While it has been accepted, copy-edited, and formatted, however, it is still undergoing proofreading and corrections by the authors. Therefore, the text may still change before the final publication. Although 'Early view' may not have all bibliographic details available, the DOI and the year of online publication can still be used to cite them. The article title, DOI, publication year, and author(s) should all be included in the citation format. Once the 'Final Version of Record' becomes available the 'Early view' will be replaced by that.