

Caries Removal by Chemomechanical (Carisolv™) vs. Rotary Drill: A Systematic Review

Viral P. Maru*, B.S. Shakuntala and C. Nagarathna

Department of Pedodontics and Preventive Dentistry, Rajarajeswari Dental College & Hospital, Mysore Road, Kumbalgodu, Bangalore 560 074 India

Abstract: *Background:* Chemomechanical caries removal is an effective alternative to the traditional rotary drilling method. The advantages of chemomechanical techniques in terms of the need for anesthesia, pain perception and patient preference are systematically reviewed and a meta-analysis of the time required for caries removal is reported. *Method:* Randomized controlled studies of comparison of chemomechanical techniques with conventional rotary drill were selected from a systematic search of standard biomedical databases, including the PubMed and Cochrane clinical trials. Non-repeated search results were screened for relevance and risk of bias assessment, followed by methodology assessment. Statistical models were applied to the outcome parameters - time required, pain perception, need of anesthesia and patient preference - extracted from the studies. *Results:* Out of the 111 non-repeated search results, 26 studies receiving a low bias score were selected for the review, and 16 randomized clinical trials of rotary and Carisolv techniques were considered for meta-analysis. Meta-analysis by fixed effect as well as random effect models indicate that Carisolv takes more time (3.65 ± 0.05 and 4.09 ± 0.29 min) than rotary drill (8.65 ± 0.09 and 8.97 ± 0.66 min) method. Advantages of reduced pain (14.67 for Carisolv vs. 6.76 for rotary drill), need for anesthesia (1.59% vs. 10.52%) outweigh the longer time requirement and make it the preferred (18.68% vs. 4.69%) method. *Conclusion:* Chemomechanical techniques stand out as a minimally invasive and preferred method based on the meta-analyses. Evaluation of pain experienced using robust methods is needed to strengthen the evidence for their use.

Keywords: Caries removal, carisolv, chemomechanical, dental caries, pain perception, rotary drill.

INTRODUCTION

Dental caries is one of the most prevalent chronic diseases, causing localized dissolution and destruction of the calcified tooth tissue. In certain cases, it may result in infection of the dental pulp [1]. There is a general agreement among clinicians that the key objective of treating carious lesions is to remove the infected layer, leaving the unaffected dentin intact [2]. It is significant to differentiate these layers to prevent the painful and unnecessary removal of sound tooth structure [2, 3].

Restorative dental treatment in children, comprising of removal of caries with conventional drill, is traumatic primarily due to fear and anxiety of children and their parents [4]. The aversion to the noise of rotary instruments and anesthesia are the key factors triggering fear and anxiety [5]. In most cases, these factors delay or avoid the uptake of dental treatment by children, leading to emergency situations. However, in such scenarios, the treatment will be even more complicated with the mandatory use of anesthesia [6]. On the other hand, there is a need for preserving tooth tissue, combined with a patient-friendly approach in every field of dentistry. The ideal strategy in restorative dentistry to ensure maximum life for the natural tooth is to preserve the sound

tissue from further damage using minimally invasive techniques [7]. Diverse techniques such as, air abrasion, atraumatic restorative technique, sono-abrasion, laser and chemomechanical methods are in use for caries removal [7-10]. Among these techniques, chemomechanical caries removal is generally considered as an effective alternative to the traditional rotary drilling method.

First chemomechanical agent for removing carious dentin was marketed under the name GK-101 in 1972 [11]. It consisted of a non-specific proteolytic agent, sodium hypochlorite in Sorensen's buffer. Even after several stages of improvement of the product and FDA approval in 1984 as Caridex, chemomechanical method gained limited acceptance due to major limitations like high cost, complexity, taste of the liquid, short life span, requirement of special equipment and large quantities of the gel, and heating. A new patented system called Carisolv™ was introduced in 1998. In contrast to Caridex, the new system uses three amino acids – lysine, leucine and glutamic acid – to enhance the effect of sodium hypochlorite on denatured collagen, and to decrease the involvement of healthy hard dental tissue, although some of the disadvantages remained [3, 12]. Papacarie™, developed in 2003, has bactericidal, bacteriostatic and anti-inflammatory characteristics due to the presence of papain, chloramines and toluidine blue salts in a thickening medium [12]. Its mechanism of action is through breaking of the partially degraded collagen molecules, thereby contributing to the degradation and elimination of the fibrin mantle formed by the carious process. The key advantages of Pa-

*Address correspondence to this author at the Department of Pedodontics and Preventive Dentistry, Rajarajeswari Dental College & Hospital, Mysore Road, Kumbalgodu, Bangalore 560 074, India; Tel: 09867220417; E-mail: viralmaru@yahoo.co.in

papacarie™ are atraumaticity, biocompatibility, antibacterial properties, non-requirement of anesthesia, and better preservation of healthy tissue while removing the compromised tissue [13].

There has been an earlier attempt at meta-analysis by Marquezan and coworkers in 2006, comparing the efficacy of chemomechanical studies with conventional drilling technique for caries removal. An observation by Marquezan *et al.* was that the studies available then did not have power of evidence generation [14-17]. Considering the various systematic, randomized clinical trials published in recent years, it is highly significant to generate clinical evidence for the use of chemomechanical caries removal methods. The objective of the present systematic review is to compare the performance of chemomechanical Carisolv™ technique with conventional rotary drill method of caries removal, in terms of (i) the pain experienced or perceived by the patients, (ii) the need for anesthesia and (iii) patients' preference of one method over the other. A secondary objective is to conduct a meta-analysis of the time for complete caries removal by Carisolv™ technique vs. conventional rotary drill method, based on randomized clinical trials published till date.

METHODOLOGY

A systematic search of original research articles, reviews and clinical studies on chemomechanical techniques of caries removal was undertaken. Published studies on the topic of comparison of techniques of caries removal were obtained based on a search of the PubMed, the Cochrane Central Register of Controlled Trials (CCTR93), Unbound Medline, Embase and Metapress databases, using the search terms 'chemomechanical', 'caries removal', 'Carisolv™', 'Papacarie™', 'clinical' and related keywords, and their combinations. The search included reports published in conference proceedings not currently listed in MEDLINE or other bibliographic databases. The search was restricted to articles published till October 2014 in the English language, by setting a language and period limitations during the search process.

All the processes of selection and evaluation of papers were performed by two expert reviewers. The selection of studies for the purpose of systematic review was performed in two stages. During the first stage of selection, as being 'relevant' or 'not relevant' (NR) to the objectives of the present review, based on the title and abstract sections. To be classified as 'relevant' and be eligible for inclusion in the second stage of selection process, a study was required to be an *in vivo* clinical study or trial of caries removal by mechanical or chemomechanical method. All types of *in vitro* studies of evaluation of the properties of caries extracted by mechanical or chemomechanical methods or restorations, comparative studies on different compositions of the gel used in chemomechanical methods, and studies in special groups of patients were classified as NR and were excluded from the second stage of selection process.

In the second stage, relevant studies were further evaluated using a Cochrane type quality assessment process based on a study of the full text of the article. Each study was assigned a risk of bias score of 'Low risk' (L) or 'High risk' (H), indicative of possible bias upon inclusion in the system-

atic review. Risk score of 'L' was assigned to randomized clinical trials (RCT), controlled clinical studies and comparative studies of rotary drill and Carisolv™ methods reporting the performance parameters of time for caries removal, need for anesthesia, pain perception and patient preference. A risk score of 'H' was given to non-comparative studies, *in vitro* studies and those using other caries removal methods and evaluating parameters other than those mentioned above. Additional stringent inclusion-exclusion criteria were applied to further shortlist studies for meta-analysis of time required for the removal of caries by conventional rotary drill vs. Carisolv™ technique.

Data on the type of study, demographic data of subjects in different groups, various performance parameters of the different techniques used for comparison, and their statistical significance were extracted from the included studies. Meta-analysis was performed based on two models: fixed effect and random effect model. Statistical measures of heterogeneity of data, Q and I² were evaluated for the two techniques. Meta-analysis of the selected data was done using SAS® (SAS Institute Inc., Cary NC USA) and Microsoft Excel (Microsoft, Inc., Redmond WA USA) statistical analysis software.

RESULTS

The search of PubMed database yielded 79 unique studies, published until October 2014. A systematic search of the CCTR93 yielded 28 trials including five 'Papacarie' trials. International Association for Dental Research (IADR) Conference proceedings had a study reported in 2005, while Unbound Medline database yielded three studies. Similar search of Metapress, Google Scholar, ClinicalTrials.gov and some journals in dental research did not yield any other unique studies. A schematic flow chart of the literature search strategy is shown in Fig. (1). Search results from these databases were pooled and a significant number of studies were identified as duplicates were excluded.




Fig. (1). Flow chart of the literature search process.

Table 1. Studies selected for systematic review, their salient features and extracted data.

No.	Author-Year	Subjects			Type of Teeth	Rotary or Mechanical method		Carisolv™		Other Measurements				Reason for non-inclusion in meta-analysis
		No. of Caries (N)	No. of Patients	Age range (y) (Mean ± SD)		N	Time (m) (Mean ± SD)	N	Time (m) (Mean ± SD)	Measure	Technique	Mech. method	Carisolv™	
1	Zinck-1988 ^a (Caridex)	114	57	NA	Previously restored with secondary decay	57	(-)3.29 ± 0.65	57	3.03 ± 0.90	Anesthesia needed		21/57 (37%)	12/57 (21%)	Older Caridex method Restored teeth removal Absolute time values not reported
2	* Ericson-1999	127	137	3-85 (35 ± 21)	Mix of 4 types of teeth	19	4.4 ± 2.2	106	10.4 ± 6.1	Anesthesia needed		9/20 (45%)	3/107 (2.8%)	-
										Degree of pain	Patient survey	1/11 'No pain' 9/11 'some pain'	58/104 'No pain' 44/104 'some pain'	
3	* Fure-2000	60	38		Primary root caries	26	4.5 ± 2.0	34	5.9 ± 2.2	Anesthesia needed		6/26 (23%)	4/34 (12%)	-
4	Maragakis-2001	32	16	7-9 (7.7 ± 0.7)	Primary molars	16	0.2 ± 0.05	16	6.85 ± 2.61	Anesthesia needed	Patient survey	16/16 (100%)	0/16 (0%)	Method of measuring time for removal of caries is different
										Patient preference	Patient survey	11/16 (68%)	5/16 (31%)	
5	Nadanovsky-2001	132	66	6-44	Permanent teeth	66	8.6 ± 3.8	66	9.2 ± 3.8	Anesthesia needed		5/66 (8%)	2/66 (3%)	Non-rotary method used, conventional spoon excavator method
										Pain perception	Patient survey	43/66 (65%) 'some pain'	21/66 (32%) 'some pain'	
6	Chaussain-Miller-2003	120	96	10-81 (35.9 ± 17.7)				94	11.1 ± 9.51	Anesthesia needed			30/94 (32%)	Not a comparative study
										Pain perception	Patient survey		68.3% 'No pain'	
7	* Kakaboura-2003	90	45	18-55		45	6.8 ± 2.8	45	12.2 ± 4.1	Anesthesia needed		40%	8%	-
										Patient preference	Patient survey	12%	88%	
8	* Rafique-2003	44	22	13-75	Contra lateral teeth	22	6.3 ± 1.3	22	5.4 ± 2.4	Anesthesia needed		100%	0%	-
										Patient acceptance	Patient survey	0%	100%	
9	* Kavvadia-2004	92	31	2-9 (4.2 ± 1)	Primary teeth	27	2.8 ± 1.9	65	8.1 ± 5.3	Anesthesia needed	Class V patient survey	4/17 (24%)	1/43 (2%) ¹¹	-
10	Fure-2004	202	170	19-85				104 98	6.7 ± 4.1 7.6 ± 4.2	Patient preference	Patient survey	19%	81%	Comparison of two different gels (Carisolv method)

(Table 1) contd....

No.	Author-Year	Subjects			Type of Teeth	Rotary or Mechanical method		Carisolv™		Other Measurements				Reason for non-inclusion in meta-analysis
		No. of Caries (N)	No. of Patients	Age range (y) (Mean ± SD)		N	Time (m) (Mean ± SD)	N	Time (m) (Mean ± SD)	Measure	Technique	Mech. method	Carisolv™	
11	* Balcuniene-2005 ^j	35	30	2.5-13	Deciduous & Permanent	30	5.9± 4.75 (1-20)	30	10.5± 4.25 (3-20)	Anesthesia needed		9/29 (31%)	1/30 (3.3%)	-
										Pain perception	Patient survey	8/29 'No pain' 15/29 'some pain'	14/30 'No pain' 7/30 'some pain'	
12	*Bergmann-2005	92	46	4-11 (8)	Maxille /Mandibular molars (deciduous)	46	3.3 ± 2.3	46	6.7 ± 2.9	Pain perception	Patient survey	18/46 'No pain' 17/46 'some pain'	26/46 'No pain' 17/46 'some pain'	-
										Patient preference	Patient interview	0%	65%	
13	* Peters-2006		50	6-11 (8.1)		24	1.34 ± 1.4	26	8.06 ± 3.13	Anesthesia needed			6/26 (23%)	-
										Pain reported			28% 'some pain'	
14	* Lozano-Chourio-2006	80	40	7-9 (7.7 ± 0.7)	Primary Molars	40	2.47 ± 1.83	40	7.51 ± 2.10	Anesthesia needed		2/40 (5%)	0/40 (0%)	-
										Pain Perception	Patient survey	24/40 'No pain' 16/40 'some pain'	33/40 'No pain' 7/40 'some pain'	
										Patient preference		11/38 (29%)	27/38 (71%)	
15	Magalhaes-2006 ^j	30			Molars	30	3.61 ± 1.17	30	6.42 ± 2.62	Knoop Hardness (KHN)	Micro-hardness Tester		Lower KHN at all distances	Non-rotary, hand excavation method
16	* Pandit-2007	150	75	6-9	Deciduous teeth	50	4.28 ± 1.67	50	8.9 ± 3.78	Pain (Mean ± SD)	VAS	4.24 ± 1.25	2.18 ± 1.12	-
										Pain (Mean ± SD)	VPS	1.44 ± 0.91	0.08 ± 0.27	
										Efficacy (Mean±SD)		0.38 ± 0.75	0.42 ± 0.76	
17	* Inglehart-2007		50	8.16		24	1.34 ± 1.4	26	8.06 ± 3.13	Perceived pain (0-4)	Operator Survey	2.42/4 (60.5%)	2.77/4 (69.25%)	-
										Satisfaction (0-5)		4.00/5 (80%)	2.62 (52.4%)	
										Pain experience (0-100)	Patient Survey	61.12 (61.12%)	69.71 (69.71%)	
										Satisfaction (0-4)		3.96/4 (99%)	3.46/4 (86.5%)	

(Table 1) contd....

No.	Author-Year	Subjects			Type of Teeth	Rotary or Mechanical method		Carisolv™		Other Measurements				Reason for non-inclusion in meta-analysis
		No. of Caries (N)	No. of Patients	Age range (y) (Mean ± SD)		N	Time (m) (Mean ± SD)	N	Time (m) (Mean ± SD)	Measure	Technique	Mech. method	Carisolv™	
18	* Hosein-2008	60	30		Mandibular molars	30	7.4 ± 3.21	30	12.19 ± 3.7	Incomplete removal	Clinical assessment	0/30 (0%)	3/30 (10%)	-
19	* Peric-2009	120	120	3-17 (8.7 ± 3.0)		60	5.2 ± 2.8	60	11.2 ± 3.3	Anesthesia needed		36/60 (60%)	7/60 (11.6%)	-
										Pain perception	Patient Survey	10/24 'No pain'	46/53 'No pain'	
												10/24 'some pain'	5/53 'some pain'	
										Incomplete removal	Clinical assessment		5/60 (8.3%)	
									Patient satisfaction	Patient Survey	28/60 (47%)	51/60 (85%)		
20	Sanjeet-2011	80	40	4-8	Primary molars	40	2.08 ± 0.38	40	5.48 ± 0.75	Pain perception	Wong Baker Faces Pain Scale	6.65 ± 1.89	1.525 ± 1.36	Non-Carisolv, Papacarie method
										Mean reduction in viable bacterial-count	Colony count	87.94%	81.12%	
21	Anegundi-2012	60	30	4-9	Primary molars	30	4.68	30	17.96	Pain perception	Patient Survey	50% 'No pain'	86.7% 'No pain'	Non-Carisolv, Papacarie method
												46.7% 'some pain'	10% 'some pain'	
										Patient preference	Patient Survey	36.7%	60%	
									Mean bacterial count	Colony count	90.33	115.5		
22	* Bohari-2012	120		5-9		30	3.45 ± 0.37	30	7.91 ± 0.72	Pain perception	FLACC Scale	2.93 ± 1.74	1.13 ± 1.25	-
										Complete removal (%)	DIAG-NODENT pen	92.9 ± 9.2	87.7 ± 6.4	
23	Matsumoto-2012	40	20	5-8	Deciduous molars	20	1.73 ± 1.3	20	2.75 ± 0.8	Pain perception		10/20 'No pain'	9/20 'No Pain'	Non-Carisolv, Papacarie method
											10 'some pain'	11 'some pain'		
24	* Goomer-2013	150	80	6-10	Primary molars	50	3.37 ± 1.1	50	14.17 ± 2.03	Pain (Mean ± SD)	VAS	77.20 ± 19.8	20.40 ± 12.28	-
										Pain (Mean ± SD)	VPS	2.72 ± 0.3	0.82 ± 0.83	
										Efficacy (removal)	Clinical assessment	0.48 ± 0.3	1.2 ± 0.83	

(Table 1) contd....

No.	Author-Year	Subjects			Type of Teeth	Rotary or Mechanical method		Carisolv™		Other Measurements				Reason for non-inclusion in meta-analysis
		No. of Caries (N)	No. of Patients	Age range (y) (Mean ± SD)		N	Time (m) (Mean ± SD)	N	Time (m) (Mean ± SD)	Measure	Technique	Mech. method	Carisolv™	
25	* Hamama-2013	32			Molars	8	4.14 ± 0.32	8	6.46 ± 1.57	Vickers Hardness, 75µm		79.16 ± 5.7	19.01 ± 2.5	-
26	Motta-2013	40	20	4-7	Primary teeth	20		20		Pain perception (Face scale) Anesthesia needed		13/20 no pain 7 some pain	18/20 no pain 2 mild pain	Non-Carisolv, Papacarie method

* Studies selected for meta-analysis of time for caries removal by the two methods
NA Not available in the article

^a Mean times adjusted for volume (residuals)

^d value inferred from the abstract

^b comparison of Carisolv™ and Papacarie™ methods

^j Mean and/or SD was calculated using the reported median, range and N values

A total of 111 non-repeating studies were evaluated on the basis of methodology and relevance to the objective of the present study, to select 56 studies during the first stage of selection. A total of 55 NR studies including *in vitro* microbiological, morphological, behavioral, radiographic, physiological, follow-up and micro-hardness studies were excluded from the present study. Among the NR studies were studies of restoration, residual caries and surface damage.

During the second stage of selection, risk of bias score was assigned to the 56 relevant studies. Twenty four studies received 'H' score for higher risk of bias while 32 studies were assigned as 'L' score. Out of the 32 low risk studies, six studies including two conference abstracts published later, and a previous meta-analysis were not selected. The final selection consisted of 26 papers and the data extracted from them are shown in Table 1.

Studies reporting the time required for the removal of caries by conventional rotary drill vs. Carisolv™ technique were shortlisted for meta-analysis. Non-randomized studies, studies with methods other than rotary and Carisolv™, and those with inadequate design (Galuscan *et al.* have not measured time and Kumar *et al.* have not used rotary method) were excluded from meta-analysis [18-21]. In addition, one study was excluded because of extreme heterogeneity of the data, probably due to difference in the method of measurement; two studies were excluded due to inadequate reporting of data (Mean and SD were not available) [22-24]. Seven other studies were excluded as non-relevant: one study for the use of hand tools for excavation, one using new gel for Carisolv™ technique, one study of Caridex system[®] one evaluated radiographically, and three studies for the use of Papacarie™ technique [25-31]. Studies by Irena *et al.* and Magalhaes *et al.* did not report the mean time and SD for removal of caries and these two parameters were calculated using the reported median and range, following the method of Hozo *et al.*, before inclusion in the meta-analysis [32-34].

Subsequent to application of the inclusion-exclusion criteria, 16 studies were selected for meta-analysis.

A summary of the studies selected for meta-analysis and their salient characteristics are shown in Table 1. Twenty six studies tabulated were diverse in terms of age groups of subjects, type of teeth caries removed *etc.* Table 1 shows the data on characteristic performance parameters of the mechanical (rotary standard technique in most of the studies) and chemomechanical (Carisolv™ in all of the selected studies, except for five studies of Papacarie™) methods of caries removal. A comparison of these parameters will be useful to determine the merits of chemomechanical method over the conventional rotary method.

Time Required for Caries Removal

Twenty two of the 26 studies tabulated, report a difference in the time for caries removal by the two methods, which is statistically significant. Based on the inclusion-exclusion criteria mentioned above, 16 studies were included for meta-analysis with respect to time for caries removal. Fig. (2A and 2B) shows the Forest plots for time taken reported by these studies for the two techniques. The data are very heterogeneous as indicated by the very high values of the I² statistic (fixed effects model) for the rotary drill (96.37%) as well as Carisolv™ (97.72%) techniques. The use of random effects model, appropriate for heterogeneous datasets, yields moderate I² values of 56.71% for the rotary drill technique, and-9.36% for the Carisolv™ technique. The effect size (or mean time for removal of caries) for the rotary and Carisolv™ techniques, determined by the random effects model are 4.10 ± 0.29 (3.52-4.67 for 95% CI) and 8.97 ± 0.66 min (7.67-10.27 for 95% CI), as indicated by the open and filled diamond symbols respectively. Similar to the observations of individual studies, these values are statistically significant (P < 0.0001), to confirm that Carisolv™ technique takes longer time for caries removal. Complete details of the meta-analysis by the two models are given in Table 2.




Fig. (2). Distribution of time for caries removal by (A) rotary drill technique (open squares, Random effects model estimate (◇, 95% CI) and (B) Carisolv™ technique (filled squares, Random effects model estimate (◆, 95% CI).

Table 2. Meta-analysis and estimation of significance levels between methods.

Study	Time required (Mean ± SD) (min)		P-Value
	Rotary or	Carisolv™	
Ericson-1999	4.4 ± 0.5	10.4 ± 0.59	< 0.0001
Fure-2000	4.5 ± 0.39	5.9 ± 0.38	0.0138
Kakaboura-2003	6.8 ± 0.42	12.2 ± 0.61	< 0.0001
Rafique-2003	6.3 ± 0.28	5.4 ± 0.51	0.1295
Kavvadia-2004	2.8 ± 0.37	8.1 ± 0.66	< 0.0001
Irena-2005	5.9 ± 0.87	10.5 ± 0.78	< 0.0001
Bergmann-2005	3.3 ± 0.34	6.7 ± 0.43	< 0.0001
Peters-2006	1.34 ± 0.29	8.06 ± 0.61	< 0.0001
Lozano-Chourio-2006	2.47 ± 0.29	7.51 ± 0.33	< 0.0001
Pandit-2007	4.28 ± 0.24	8.9 ± 0.53	< 0.0001
Inglehart-2007	1.34 ± 0.29	8.06 ± 0.61	< 0.0001
Hosein-2008	7.4 ± 0.59	12.19 ± 0.68	< 0.0001
Peric-2009	5.2 ± 0.36	11.2 ± 0.43	< 0.0001
Bohari-2012	3.45 ± 0.07	7.91 ± 0.13	< 0.0001
Goomer-2013	3.37 ± 0.16	14.17 ± 0.29	< 0.0001
Hamama-2013	4.14 ± 0.11	6.46 ± 0.56	0.0011
Meta Analysis			
Fixed Effect Model	3.65 ± 0.05	8.65 ± 0.09	< 0.0001
	I ² = 96.37 %	I ² = 97.72 %	
Random Effect Model	4.09 ± 0.29	8.97 ± 0.66	< 0.0001
	I ² = 56.71 %	I ² = - 9.36 %	




Fig. (3). Distribution of percentage of patients who perceived ‘No pain’ (A) or ‘some pain’ (B) during caries removal by rotary drill technique (open squares, mean ----, weighted mean ◇) and chemomechanical techniques (filled squares, mean —, weighted mean ◆).

Evaluation of Pain

Clinical measures of pain perceived, need for anesthesia and patient preference have been analyzed on the basis of weighted average of the data from selected studies. Fifteen studies have reported measure of degree of pain experienced by the patient during the procedure of caries removal. These results are consistent with the fact that rotary drilling method is more painful compared to the chemomechanical method. Pain experienced during the procedure is expressed in the patient surveys by the two options: ‘no pain’ or ‘some pain’ experienced by the patient. Although other options were also employed in some studies, majority of the subjects have used these options and the difference between the techniques is most pronounced with these options. These two patient responses have been analyzed separately, as shown in Fig. (3A and 3B). The simple mean of percentage of ‘no pain’ and ‘some pain’ responses for the rotary drill technique is shown by dotted lines and those for Carisolv™ are shown by solid lines in Fig. (3A and 3B). The weighted average for these responses in each case are represented by the open diamonds and filled diamonds for rotary and Carisolv™ techniques respectively. The case of ‘no pain’ responses, simple average value is very high for Carisolv™ (30.44% for rotary compared to 64.32% for the chemomechanical method), implying that more subjects did not experience pain with Carisolv™ technique. However, the weighted average for rotary technique is only slightly smaller (16.28) than that for Carisolv™ technique (17.12), indicating that there is no significant difference between the techniques. The difference between the techniques in terms of pain perceived is unequivocally demonstrated by considering the positive choice of ‘some pain’ experienced by the patient. A reverse of the earlier trend is true for ‘some pain’ responses – higher

weighted average value for rotary technique (14.67) reflects that many subjects experienced pain with rotary drill and only a smaller number (6.76) experienced pain during Carisolv™ extraction.

Need for Anesthesia

Requirement of anesthesia during caries removal by the patients has been reported in 12 out of 27 selected studies. Majority of subjects assigned to rotary method opted for anesthesia as indicated by their percentage ranging from 23 to 60%, and occasionally to 100% of them have been treated under anesthesia [22, 35]. On the other hand, in most of the Carisolv™ studies, 0 to 12% of subjects opted for anesthesia and only in two studies, their proportion reached 20 to 30% [19, 36]. Distribution of subjects opting for anesthesia in various studies is shown in Fig. (4). The weighted mean of subjects for rotary technique is clearly higher (10.52%) than those opting for anesthesia during chemomechanical caries removal (1.59%). Less than 10% of the group population needed anesthesia in most of the chemomechanical studies, while >30% subjects are treated under anesthesia for the rotary technique.

Patient Preference

Patient surveys in the form of written questionnaire or interview by clinicians have been conducted after the caries removal process. Survey has been useful to quantify many subjective experiences about the process the patients have just undergone. Patients’ preference to one of these methods over the other may sum up these experiences. Patient preferences recorded in seven studies have been plotted as shown in Fig. (5), and analyzed using the weighted average of the

percentage of subjects preferring each technique. In all studies, except one, more than 50% subjects preferred Carisolv™ technique to rotary and in two studies, none of the subjects preferred the rotary drill technique [22, 35, 37]. The weighted average of 18.68% for Carisolv™, which is higher than that for rotary drill (4.69%), confirms the trend.




Fig. (4). Distribution of percentage of patients who needed anesthesia during rotary drill technique (open squares, mean -----, weighted mean ◇) and chemomechanical techniques (filled squares, mean —, weighted mean ◆).




Fig. (5). Distribution of patient preference to rotary drill technique (open squares, mean -----, weighted mean ◇) and chemomechanical techniques (filled squares, mean —, weighted mean ◆).

DISCUSSION

The present meta-analysis of the comparative clinical studies of the mechanical and chemomechanical techniques of caries removal has shown that the time required for both

methods was significantly different. Time for removal has been taken as the key parameter for meta-analysis, since there is significant heterogeneity in the reporting of other treatment parameters. Measures of pain perception, requirement of anesthesia and patient preference have been evaluated in fewer studies and by different scales and techniques, and hence, are not amenable for meta-analysis. The time required for complete caries removal is significantly more than that for conventional rotary drill, but the other advantages in terms of reduced pain and need for anesthesia outweigh the longer duration.

Although the time required for complete removal of caries by mechanical rotary drill as well as the chemomechanical Carisolv™ or Papacarie™ methods has been reported in 22 studies (Table 1), all these studies could not be included in the meta-analysis due to various reasons, explained above. It is clear from the meta-analysis that chemomechanical Carisolv™ technique takes longer duration for caries removal. Apart from the Carisolv™ studies considered for meta-analysis, few studies have compared the other chemomechanical Papacarie™ with rotary and / or Carisolv™ techniques. In a study by Kumar *et al.*, Papacarie™ technique took a marginally less mean time of 10.48 ± 2.96 min. for caries removal compared to 11.67 ± 3.25 min. (P>0.05) for Carisolv™ technique [21]. Similar results have been obtained by Kotb *et al.* [29] and Kochhar *et al.* [38]. Matsumoto *et al.* did not find significant difference between the time taken by Papacarie™ Duo and low speed bur techniques [31].

Among the patient surveys of pain perception, only two of the most frequent options selected by the patients are used in the present study for further analysis. Responses for the complete absence of pain indicated by ‘no pain’, and presence of some tolerable pain indicated by ‘some pain’ have been consistently used by significantly higher number of patients for chemomechanical methods than for the rotary method. For rotary technique, the number of ‘no pain’ responses is usually below 30%, but in two studies, these responses were as high as 40-60%. [37, 39]. In the case of chemomechanical technique, ‘no pain’ responses were consistently above 50% and in two cases, as high as 75-85% [39, 40]. When the second option of ‘some pain’ is considered, the trend becomes clear in that responses for the Rotary range from 40-65% while those for Carisolv™ method ranges from 15-30%.

Inglehart *et al.* have used both operator and patient surveys for the evaluation of pain experienced during caries removal by the two methods [41]. The operator perceived pain was scored on a scale of 0-4, whereas patients rated their pain experience on scale of 0-100. In their study, average pain experienced during the rotary technique was only slightly less (60%) than the chemomechanical method (69%). Interestingly, both the operator and patients arrive at similar values for the pain experienced by the two methods, in spite of the difference in evaluation methods.

Evaluation of pain using structured pain scales by trained clinicians is expected to yield a better estimate of pain experience, instead of patient surveys, as individual tolerance to pain may influence patient response to the survey. Pain evaluation has been carried out using Visual Analog Scale

(VAS) and Verbal Pain Scale (VPS) by Pandit *et al.* and Goomer *et al.*, using Wong Baker Faces Pain Scale by Sanjeet *et al.*, and using the FLACC (Face, Leg, Activity, Cry, Consolability) scale by Bohari *et al.* [30, 42-44]. Once again, pain experienced during Carisolv™ method ranks consistently lower than rotary drill method, irrespective of the scale used. However, when mean values of pain by VAS scale across studies are compared, Goomer *et al.* report a very high 77.20 ± 19.8 compared to 4.24 ± 1.25 reported by Pandit *et al.* [42-45]. Overall, pain experience by the rotary drill method is clearly higher than the Carisolv™ method.

Thirteen of the selected studies have determined the number (or percentage) of patient who were given or opted for anesthesia during caries removal by the two techniques. Of these, Chousian-Miller *et al.* [19] did not report a value for the rotary drill method as it is a non-comparative study of Carisolv™ method alone. The extent of pain experienced and/or the fear of patients, especially children, for the rotary drill method is indicated by the percentage of them opting for anesthesia for the procedure. Out of the 11 studies reporting this measure, in nine studies >25% of the patients needed anesthesia and in two of the studies, 100% of the patients were treated under anesthesia. On the other hand, requirement of anesthesia for the chemomechanical method is much lower, as indicated in Table 1. In twelve studies (out of the 13) where Carisolv™ was used, <25% of the subjects needed anesthesia and in the remaining study also, the number was a moderate 32%. None of the subjects required anesthesia in studies by Maragakis *et al.*, Rafique *et al.* and Lozano-Chourio *et al.*, while <5% were given anesthesia in studies by Ericson *et al.*, Nadanovsky *et al.*, Kavvadia *et al.* and Irena *et al.* [22, 25, 32, 35, 39, 46-49]. All of these observations go a long way to establish Carisolv™ as a mild and non-traumatic technique for caries removal, especially in children.

In eight of the selected studies, subjects answered a questionnaire or were interviewed after the treatment, about their experience with the two techniques and their preference to one technique over the other. Patient 'preference' has been differentiated from 'patient acceptance' reported by Rafique *et al.* and 'patient satisfaction' reported in two other studies. [35, 40, 41] In the present analysis, only patient preference measured as a percentage of that specific group population was included. The distribution of patient preferences for the chemomechanical over the rotary drill technique is indicated in Fig. (5). Chemomechanical method clearly stands out as the most preferred, based on the higher weighted average value compared to that for rotary drill technique.

CONCLUSION

In summary, the present systematic review and meta-analysis have established that Carisolv technique has relatively superior performance in terms of the need for anesthesia, pain experience by the patients and preference over other methods, in spite of taking longer time for caries removal. Many randomized controlled studies of chemomechanical techniques have been published over the last decade. Time for removal is the only parameter that could be compared by the current meta-analysis. Comparison of other important clinical parameters of pain perception or pain experienced by

robust methods is essential to strengthen the evidence for the use of chemomechanical techniques [50]. In comparison to the studies considered for meta-analysis by Marquezan *et al.*, the quality of studies has improved in recent times, but sample sizes are very small in many studies (especially Hamama *et al.*). [17, 45] In spite of these limitations, chemomechanical technique stands out as a minimally invasive and preferred method compared to conventional rotary drill, based on the analyses done here. Chemomechanical caries removal techniques, being non-traumatic and least-injurious to healthy tooth structure, hold excellent promise for future dental practice [51].

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Lundeen TF, Roberson TM. In: Roberson TM, Heymann HO, Swift EJ, Ed. Sturdevant's Art and science of operative dentistry. St Louis: Mosby Co 2002; pp. 60-128.
- [2] Fusayama T, Kurosaki N. Structure and removal of carious dentin. *Int Dent J* 1972; 22: 401-11.
- [3] Beeley JA, Yip HK, Stevenson AG. Chemomechanical caries removal: a review of techniques and latest developments. *Br Dent J* 2000; 188: 427-30.
- [4] Scott S, Hirschman P, Scroder K. Historical antecedents of dental anxiety. *J Am Dent Assoc* 1984; 108: 42-5.
- [5] Ayer A, Domoto K, Gale N, Joy D, Melamed G. Overcoming dental fear: strategies for its prevention and management. *J Am Dent Assoc* 1983; 107: 18-27.
- [6] Michelle M, Luis F. Chemomechanical caries removal: current evidences. *RBO* 2005; 62: 125-9.
- [7] Banerjee A, Kidd E, Watson T. Scanning electron microscope observations of human dentin after mechanical caries excavation. *J Dent* 2000; 28: 179-86.
- [8] Myers GE. The air abrasive technique. *Br Dent J* 1954; 97: 291-5.
- [9] Frencken JE, Pilot T, Sangpaison Y, Phantumvanit P. Atraumatic restorative treatment: rationale, technique and development. *J Public Health Dent* 1996; 56: 135-40.
- [10] Keller U, Hibst R, Geurtsen W, *et al.* Erbium: YAG laser application in caries therapy. Evaluation of patient perception and acceptance. *J Dent* 1998; 26: 649-56.
- [11] Goldman M, Kronman JH. A preliminary report on a chemomechanical means of removing caries. *J Am Dent Assoc* 1976; 93: 1149-53.
- [12] Silva LR. Papacarie : A new system for the chemomechanical caries removal: case report. *Rev Paul Odontol* 2004; 16: 4-8.
- [13] Douglas C, Ana F, Franciele O, Eloisa B, Bruno C. Effect of Carisolv and Papacarie on the resin dentin bond strength in sound and caries affected primary molars. *Braz J Oral Sci* 2010; 9: 25-9.
- [14] Montedori A, Abraha I, Orso M, D'Errico PG, Pagano S, Lombardo G. Lasers for caries removal in deciduous and permanent teeth (Protocol). *Cochrane Database Syst Rev* 2012; Issue 11: Art. No.: CD010229. DOI: 10.1002/14651858.CD010229.
- [15] Ricketts D, Lamont T, Innes NPT, Kidd E, Clarkson JE. Operative caries management in adults and children. *Cochr Database Syst Rev* 2013; Issue 3: Art. No.: CD003808. DOI: 10.1002/14651858.CD003808.pub3.
- [16] Innes NPT, Ricketts D, Evans DJP. Preformed metal crowns for decayed primary molar teeth. *Cochr Database System Rev* 2007; Issue 1: Art. No.: CD005512. DOI: 10.1002/14651858.CD005512.pub2.
- [17] Marquezan M, Faraco Junior IM, Feldens CA, Tovo MF, Ottoni AB. Evaluation of the methodologies used in clinical trials and

- effectiveness of chemo-mechanical caries removal with Carisolv™. *Braz Oral Res* 2006; 20: 364-71.
- [18] Munshi AK, Hegde AM, Shetty PK. Clinical evaluation of Carisolv in the chemico- mechanical removal of carious dentin. *J Clin Pediatr Dent* 2001; 26: 49-54.
- [19] Chaussain-Miller C, Decup F, Domejean-Orliaguet S, *et al.* Clinical evaluation of the Carisolv chemomechanical caries removal technique according to the site/stage concept, a revised caries classification system. *Clin Oral Investig* 2003; 7: 32-7.
- [20] Galuscan A, Jumanca D, Podariu A, Ardelean L, Rusu LC. Evaluation by fluorescent light of chemomechanical treatment of caries removal using Carisolv™. *Rev Chim (Bucharest)* 2012; 63: 949-52.
- [21] Kumar J, Nayak M, Prasad KL, Gupta N. A comparative study of the clinical efficiency of chemomechanical caries removal using Carisolv and Papacarie: a papain gel. *Indian J Dent Res* 2012; 23: 697.
- [22] Maragakis GM, Hahn P, Hellwig E. Clinical evaluation of chemomechanical caries removal in primary molars and its acceptance by patients. *Caries Res* 2001; 35: 205-10.
- [23] Lozano MA, Zambrano O, Gonzalez H. Clinical randomized controlled trial of chemomechanical caries removal (Carisolv™). In: *Proceedings of the 83rd Intl Assoc Dent Res (IADR) Conference 2005*: Baltimore, MD.
- [24] Aneundi RT, Patil SB, Tegginmani V, Shetty SD. A comparative microbiological study to assess caries excavation by conventional rotary method and a chemomechanical method. *Contemp Clin Dent* 2012; 3: 388-92.
- [25] Nadanovsky P, Cohen Carneiro F, Souza de Mello F. Removal of caries using only hand instruments: a comparison of mechanical and chemo-mechanical methods. *Caries Res* 2001; 35: 384-9.
- [26] Fure S, Lingstrom P. Evaluation of the chemomechanical removal of dentin caries *in vivo* with a new modified Carisolv™ gel. *Clin Oral Investig* 2004; 8: 139-44.
- [27] Zinck JH, McInnes-Ledoux P, Capdeboscq C, Weinberg R. Chemomechanical caries removal – a clinical evaluation. *J Oral Rehab* 1988; 15: 23-33.
- [28] Figuerado MC, Machado CV, Castro ME. Clinical and radiographic evaluation of deciduous teeth submitted to Carisolv™ for caries removal. *Intl J Pediatr Dent* 2003; 13(Suppl-1): 5.
- [29] Kotb R, Abdella A, Kateb M, Ahmed A. Clinical evaluation of Papacarie™ in primary teeth. *J Clin Pediatr Dent* 2009; 34: 117-24.
- [30] Singh S, Jawa Singh D, Jaidka S, Somani R. Comparative clinical evaluation of chemomechanical caries removal agent Papacarie® with conventional method among rural population in India - *in vivo* study. *Braz J Oral Sci* 2011; 10: 193-8.
- [31] Matsumoto SB, Motta LJ, Alfaya TA, Guedes CC, Fernandes KS, Bussadori SK. Assessment of chemomechanical removal of carious lesions using Papacarie Duo™: Randomized longitudinal clinical trial. *Ind J Dent Res* 2013; 24: 488-92.
- [32] Balciuniene I, Sabalaite R, Juskiene I. Chemomechanical caries removal for children. *Stomatol Baltic Dent Maxillofac J* 2005; 7: 40-4.
- [33] Magalhaes CS, Moreira AN, Campos WR, Rossi FM, Castilho GA, Ferreira RC. Effectiveness and efficiency of chemomechanical carious dentin removal. *Braz Dent* 2006; 17: 63-7.
- [34] Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range and the size of a sample. *BMC Med Res Methodol* 2005; 5: 13.
- [35] Rafique S, Fiske J, Banerjee A. Clinical trial of an air-abrasion/chemomechanical operative procedure for the restorative treatment of dental patients. *Caries Res* 2003; 37: 360-4.
- [36] Peters MC, Flamenbaum MH, Eboda NM, Feigal RJ, Inglehart MR. Chemomechanical caries removal in children: efficacy and efficiency. *J Amer Dent Assoc* 2006; 137: 1658-66.
- [37] Bergmann J, Leitão J, Kultje C, Bergmann D, Clode MJ. Removing dentine caries in deciduous teeth with Carisolv: a randomised, controlled, prospective study with six-month follow-up, comparing chemomechanical treatment with drilling. *Oral Health Prev Dent* 2005; 3: 105-11.
- [38] Kochhar GK, Srivastava N, Pandit I, Gugnani N, Gupta M. An evaluation of different caries removal techniques in primary teeth: a comparative clinical study. *J Clin Pediatr Dent* 2011; 36: 5-10.
- [39] Lozano-Chourio MA, Zambrano O, Gonzalez H, Quero M. Clinical randomized controlled trial of chemomechanical caries removal (Carisolv™). *Intl J Pediatr Dent* 2006; 16: 161-7.
- [40] Peric T, Markovic D, Petrovic B. Clinical evaluation of a chemomechanical method for caries removal in children and adolescents. *Acta Odontol Scand* 2009; 67: 277-83.
- [41] Inglehart MR, Peters MC, Flamenbaum MH, Eboda NN, Feigal RJ. Chemomechanical caries removal in children: an operator's and pediatric patients' responses. *J Am Dent Assoc* 2007; 138: 47-55.
- [42] Pandit IK, Srivastava N, Gugnani N, Gupta M, Verma L. Various methods of caries removal in children: A comparative clinical study. *J Ind Soc Pedod Prev Dent* 2007; 93-6.
- [43] Goomer P, Jain RL, Kaur H, Sood R. Comparison of the efficacy of chemomechanical caries removal with conventional methods: A clinical study. *J Int Oral Health* 2013; 5: 42-7.
- [44] Bohari MR, Chunawalla YK, Ahmed BMN. Clinical evaluation of caries removal in primary teeth using conventional, chemomechanical and laser technique: An *in vivo* study. *J Contemp Dent Pract* 2012; 13: 40-7.
- [45] Hamama HH, Yiu CKY, Burrow MF, King NM. Chemical, morphological and microhardness changes of dentine after chemomechanical caries removal. *Aust Dent J* 2013; 58: 1-10.
- [46] Ericson D, Zimmerman M, Raber H, Gotrick B, Bornstein R, Thorell J. Clinical evaluation of efficacy and safety of a new method for chemo-mechanical removal of caries. *Caries Res* 1999; 33: 171-7.
- [47] Fure S, Lingström P, Birkhed D. Evaluation of Carisolv™ for the chemo-mechanical removal of primary root caries *in vivo*. *Caries Res* 2000; 34: 275-80.
- [48] Kakaboura A, Masouras C, Staikou O, Vougiouklakis G. Comparative clinical study on the Carisolv caries removal method. *Quintessence Int* 2003; 34: 269-71.
- [49] Kavvadia K, Karagianni V, Polychronopoulou A, Papagiannoulis L. Primary teeth caries removal using the Carisolv™ chemomechanical method: A clinical trial. *Pediatr Dent* 2004; 26(1): 23-8.
- [50] Hosein T, Hasan A. Efficacy of chemo-mechanical caries removal with Carisolv. *J Coll Physic Surg Pak* 2008; 18: 222-5.
- [51] Motta LJ, Bussadori SK, Campanelli AP, *et al.* Pain during removal of carious lesions in children: A randomized controlled clinical trial. *Intl J Dent* 2013; Article ID 896381. doi.org/10.1155/2013/896381

Received: December 10, 2014

Revised: September 29, 2015

Accepted: October 27, 2015

© Maru *et al.*; Licensee Bentham Open.

This is an open access articles licensed under the terms of the Creative Commons Attribution-Non-Commercial 4.0 International Public License (CC BY-NC 4.0) (<https://creativecommons.org/licenses/by-nc/4.0/legalcode>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided that the work is properly cited.