



The Open Dentistry Journal

Content list available at: <https://opendentistryjournal.com>



RESEARCH ARTICLE

Anti-Quorum Sensing Effect of *Salvadora Persica* Against *Enterococcus faecalis* (ATCC 29212)

Ehsan Kadhim^{1,*}, Bushra Amin² and Bassam Amin¹

¹Department of Conservative, Hawler Medical Univeristy, Erbil, Iraq

²Department of Biology, Salahaddin University, Erbil, Iraq

Abstract:

Introduction:

Quorum Sensing (QS) is a mechanism many bacteria use to manage their cooperative activities and physiological functions. The Fsr system in *Enterococcus faecalis* (ATCC 29,212) is an example of quorum sensing with a cell density-dependent two-component regulatory system mechanism. Several publications have shown that the Fsr system and proteases independently contribute to *E. faecalis* pathogenicity in various infection models.

Objectives:

There is currently no published research to determine the exact molecular ability of *Salvadora persica* on quorum-sensing genes. Therefore, this study aimed to determine the plant extracts that inhibit the expression of the quorum-sensing gene (*FsrC*).

Methods:

Different fractions of *Salvadora persica* were obtained using different solvents, including standard hexane, chloroform, ethyl acetate, n-butanol, ethyl alcohol, and water which are expressed as fractions 1,2,3,4,5 and 6, respectively. Antibacterial activity assay of different plant extracts (*S. persica*) was determined by minimum inhibitory concentration (MIC). Finally, the relative expression of the quorum-sensing (QS) gene was evaluated using a One-step quantitative RT-PCR PrimeScript™ RT-PCR Kit.

Results:

All fractions of *S. persica* showed antimicrobial activity. However, ethyl acetate- *S. persica* inhibited the growth of *E. faecalis* (ATCC 29,212) at the lowest concentration, which was 20mg/ml and the highest concentration inhibited the growth of *E. faecalis* (ATCC 29,212) was 60mg/ml (chloroform- *S. persica*). Furthermore, the highest change fold value of (4.99) was recorded in treated *E. faecalis* (ATCC 29,212) with fraction 1 (hexane).

Conclusion:

Overall, *S. persica* showed antimicrobial activity against *E. faecalis* (ATCC 29,212). However, more studies are required to investigate the effect of different plant extracts on quorum-sensing genes of *Enterococcus faecalis*.

Keywords: Antimicrobial, Expression, Endodontic, Siwak, Oral, Quorum sensing.

Article History

Received: February 07, 2022

Revised: February 15, 2022

Accepted: March 18, 2022

1. INTRODUCTION

The most prevalent enterococcal species found in clinical and dietary samples are *Enterococcus faecalis* and *Enterococcus faecium*, which are responsible for around 90%

and 10% of human nosocomial infections, respectively [1 - 6].

Quorum sensing (QS) is a mechanism used by many bacteria to manage their cooperative activities and physiological functions [7, 8]. Quorum sensing (QS) is a population-based control based on self-secreted molecules known as auto-inducers (AI); they can accumulate in the environment and change the expression of many genes [9]. Pathogenicity, biofilm formation, motility, sporulation,

* Address of Correspondence to this author at the Department of Conservative, Hawler medical Univeristy, Erbil, Iraq; Tel: 07505903637; E-mail: doctorihsanmansoor@gmail.com

conjugal plasmid transfer, bioluminescence, antibiotic resistance, and antibiotic synthesis are all regulated by quorum sensing (QS) [10, 11]. Generally, the QS systems in bacteria are categorised into at least three classes, including LuxI/LuxR-type QS in Gram-negative bacteria, oligopeptides two-component-type QS in Gram-positive bacteria, and (3) luxS-encoded autoinducer 2 (AI-2) QS in both Gram-negative and Gram-positive bacteria [11, 12]. The Fsr system in *E. faecalis* is an example of QS with a cell density-dependent two-component regulatory system mechanism [13]. The *fsr* locus is 2.8 kb in size and comprises four genes: *fsrA*, *fsrB*, *fsrD*, and *fsrC* [8, 14] and the Fsr system is required for four genes including *gelE*, *sprE*, *ef1097*, and *ef1097b* [15]. *FsrC* encodes the transmembrane histidine protein kinase FsrC, the sensor-transmitter of the *fsr* operon [14]. However, the Fsr system has been identified as a quorum-sensing locus that responds to the extracellular accumulation of a peptide lactone encoded at the C terminus of the FsrB protein [14]. The FsrC histidine kinase is presumably triggered by the accumulation of the peptide in the extracellular space, resulting in the activation of the response regulator and transcription factor FsrA [16].

Several publications have shown that both Fsr and proteases independently contribute to *E. faecalis* pathogenicity in various infection models [17 - 22]. The specific mechanisms through which Fsr and its regulatory proteases contribute to infection toxicity are unknown [15]. As mentioned previously, the accumulation of these molecules changes the expression of many genes, including the genes responsible for forming biofilm [23]. A previous study revealed that quorum-sensing is significant in biofilms and may result in dispersion [24]. Antibiotics are used to treat various infectious illnesses [25]. However, the overuse of antibiotics resulted in drug-resistant bacterial strains, including (*E. faecalis*) [26]. There is a great demand for new agents and natural compounds because of low effectiveness due to the development of resistance, host toxicity, and various side effects of currently used therapeutics [27]. As a result, blocking bacterial quorum-sensing (QS) systems have gotten a lot of attention since it has the potential to reduce bacterial virulence without causing bactericidal pressure, which can lead to drug resistance. Actinobacteria continue to produce novel molecules with various bioactive properties, such as antibacterial, anticancer, and antihelminthic properties as well as the emergence of integrative metabolomic and genomic analysis has provided new hope for natural product researchers [28]. Recent study confirmed that Pantoea Natural Product 3 (PNP-3) is a broad-spectrum antimicrobial that is effective against a variety of multidrug-resistant bacteria [29]. Discovering novel natural product with antimicrobial activity is critical as drug-resistant bacteria have few therapeutic options.

S. persica (Miswak-Siwak) is a Middle Eastern native bushy plant whose parts have been used as a natural toothbrush and mouth rinse from time immemorial [30, 31]. However, there is currently no published research to determine the exact molecular capability of *S. persica* to QS genes. Therefore, this study aimed to determine the plant extracts (*S. persica*) that inhibit the expression of the QS gene (*FsrC*).

2. MATERIALS AND METHODS

2.1. Bacterial Culture

Enterococcus faecalis (ATCC 29,212) were used in all experiments obtained from the American Type Culture Collection (ATCC), cultured out twice on BHI (Scharlau Chemie S.A., Barcelona, Spain) agar plates for 24 hours at 37°C and checked for purity by Gram stain and colony morphology [32].

2.2. Plant Extracts

Different solvents were used to extract active substances from plants; the solvents are selected primarily based on the polarity of the solute, from low polar to high polar, as follows: Hexane – Chloroform – Ethylacetate – Butanol – Ethanol – Water [33]. All solvents used for the extraction process were used as blanks and controls in all experiments throughout the study to exclude the effect of the solvents. The resultant liquid was filtered through a filter paper into a round-bottom flask and concentrated using a Buchi R-200 rotary evaporator, thus removing all the solvents and resulting in a viscous crude plant extract. The crude extracts were stored at - 20°C and freshly dissolved in 10% DMSO before use.

The mixture was prepared from 50g of dried Miswak (*S. persica*), and 100ml of Hexane put in an ultrasonic bath (1.5 hours at 37°C). Then the mixture was filtrated to get extract part one (fraction 1) solvent removed using the rotary evaporator. The residue then dried at room temperature and was used for the next solvent extraction. The same technique was used for the other solvents to get the different extract fractions (fraction 2 from chloroform, fraction 3 from ethyl acetate, fraction 4 from butanol, fraction 5 from ethanol, and fraction 6 from water).

2.3. Determination of Minimum Inhibitory Concentration (MIC)

The minimum inhibitory concentration (MIC) for antibacterial activity assay of different plant extracts was determined by serial dilution, as described by Eloff [34], according to the Clinical and Laboratory Standards Institute (CLSI) standard protocol [35]. In 1.5 ml microcentrifuge tubes (Eppendorf), stock solutions of the respective plant extracts were prepared by dissolving the dry plant extract in dimethylsulphoxide (DMSO) to a final concentration of 100mg/ml. The serial dilutions from the stock solution were made in 2ml-96-well microplates to the concentration of (5, 10, 20, 25, 30, 35, 40, 50, 60, 70 and 100 mg/ml). Mueller-Hinton broth was used to obtain an activated bacterial suspension with a concentration of 1.5×10^8 CFU/ml (0.5 MacFarland). 100µl of bacterial suspension was transferred into each well and inoculated at 37°C for 24 hours to activate. The growth control and sterile control were used for this assay; growth control contained only bacterial suspension, and sterile control contained only plant extracts. The effect of solvents on the bacterial growth was not recorded as a stock solution prepared from the dry crude extract and 10% DMSO solution. MIC was determined by reading the absorbency at wavelength 590 nm, and the lowest concentration of each extract displaying no

visible growth was recorded as the minimum inhibitory concentration. All samples were tested in triplicate.

2.4. RNA Extraction and Preparation

Total RNA was isolated by RNxplus kit (SinaClon, Tehran, Iran) from untreated and treated samples *E. faecalis* (ATCC 29,212). The RNA from treated samples was extracted after determining the minimum inhibitory concentration (MIC). The bacterial culture with the highest plant concentration that showed visible growth (a concentration before MIC) was selected for the RNA extract and considered a treated sample. Briefly, bacterial cells were dissolved in 1ml ice-cold RNxplus solution, gently shaken after adding 200 ml chloroform, and then kept on ice for 15 minutes, followed by cold centrifugation at 12000rpm for 15 minutes. Wqualvolume of isopropanol was added, followed by cold centrifugation at 12000 rpm for 15 minutes. The supernatant was discarded, 1ml Ethanol (75%) was added, and then centrifuged at 7500 rpm for 8 minutes. The supernatant was removed, and the pellet was left to dry at room temperature. Finally, it dissolved in 50µl DEPC (Diethylpyrocarbonate) (SinaClon) water. RNase-free DNase was used to eliminate DNA from the RNA fraction. Additionally, Nanodrop (BioTech, USA) was used to determine the quality of isolated RNA and immediately quantified by RT-PCR.

2.5. Quantitative RT-PCR Assay

The nucleic acid sequence of the QS gene (*FsrC*) was obtained from *Enterococcus faecalis* (ATCC 29,212). Forward

primer (GCTTATTGGAAGAACAACGTATCAA) and reverse primer (CGAAACATCGCTAGCTCTTCGT) were selected to evaluate the expression of the target QS gene [36]. Following the manufacturer's instructions, the relative expression of the QS gene (*FsrC*) was evaluated using a One-step quantitative RT-PCR PrimeScript™ RT-PCR Kit. The total volume was 20 µl, which contained 1 µl of each primer sequence, 2 µl of 2X reaction buffer, 1 µl of the enzyme solution, and 3 µl of extracted RNA. The reaction was carried out at 50°C for 20 minutes (cDNA synthesis), with the initial activation step at 94°C for 5 minutes, followed by 40 cycles of denaturation at 94°C for 25 seconds, annealing at 63°C for 60 seconds, and extension at 68°C for 60 seconds. All results were normalized by untreated samples. The effect of plant extracts on the quorum-sensing gene was determined using comparative critical threshold (Ct) real-time PCR. The following formula ($\Delta Cq = Cq \text{ treated bacteria} - Cq \text{ control bacteria}$) and $normalize = 2^{-\Delta Cq}$ are used to determine the change fold and normalize change value. All experiments were performed twice.

3. RESULTS

3.1. Minimum Inhibitory Concentration (MIC)

All fractions of *S. persica* showed antimicrobial activity, as shown in Table 1 and Fig. (1). However, ethyl acetate- *S. persica* inhibited the growth of *E. faecalis* at the lowest concentration, which was 20mg/ml, and the highest concentration, which was 60mg/ml (chloroform- *S. persica*).




Fig. (1). Different fractions and concentrations of *S. persica* and the growth of *E. faecalis* (ATCC 29,212).

Table 1. The growth of *E. faecalis* (ATCC 29,212) and different fractions of *S. persica*

<i>S. persica</i>	5mg/ml	10mg/ml	20mg/ml	25mg/ml	30mg/ml	35mg/ml	40mg/ml	50mg/ml	60mg/ml	70mg/ml	100mg/ml
Fraction 1	0.955	0.702	0.131	0	0	0	0	0	0	0	0
Fraction 2	2.343	1.911	1.802	1.408	0.777	0.563	0.183	0.142	0	0	0

(Table 1) contd....

<i>S. persica</i>	5mg/ml	10mg/ml	20mg/ml	25mg/ml	30mg/ml	35mg/ml	40mg/ml	50mg/ml	60mg/ml	70mg/ml	100mg/ml
Fraction 3	0.981	0.546	0	0	0	0	0	0	0	0	0
Fraction 4	0.857	0.220	0.181	0	0	0	0	0	0	0	0
Fraction 5	1.793	1.734	0.753	0	0	0	0	0	0	0	0
Fraction 6	0.732	0.158	0.134	0	0	0	0	0	0	0	0

Table 2. The effects of different fractions of *S. persica* on *FsrC* gene expression.

Parameter	fraction 1	fraction 2	fraction 3	fraction 4	fraction 5	fraction 6
normalize change value	0.03147	0.69255	0.31208	1.01396	1.00000	1.01396
Change fold	4.99	0.53	1.68	-0.02	0.00	-0.02

3.2. Effect of *S. Persica* on the Expression of QS *FsrC* Gene

The molecular effect of *S. persica* extract against *E. faecalis* (ATCC 29,212) was determined by evaluating the relative expression levels of *FsrC* genes. For this purpose, Real-time quantitative polymerase chain reaction (RT-qPCR) was used. As shown in Table 2 and Figs. (2 and 3), fractions 4,

5, and 6 of *S. persica* showed no effect on the level of *FsrC* expression, while the expression of the same gene decreased in bacterial samples that were treated with fractions 1, 2 and 3 of *S. persica*. Furthermore, the highest change fold value of (4.99) was recorded in treated *E. faecalis* (ATCC 29,212) with fraction 1, as shown in Fig. (3).

4. DISCUSSION

Currently, *E. faecalis* is widely known as one of the most common bacteria in the root canal [12, 37]. On the other hand, several investigations reported that both Fsr and proteases independently contribute to *E. faecalis* pathogenicity in various infection models [17 - 20, 22]. The QS system encoded by the *fsr* gene cluster on the *E. faecalis* chromosome regulates the production of two pathogenicity-related proteases, gelatinase (*GelE*) and serine protease (*SprE*) [25]. Moreover, deletion of *fsrA*, *fsrB*, or *fsrC* inhibits the expression of *gelE* and *sprE* in wild-type *E. faecalis* strain OG1RF; nonetheless, both were strongly expressed in wild-type *E. faecalis* strain OG1RF [38].

The rising incidence of antibiotic resistance by pathogenic microorganisms is an alarming issue in treating diseases. There is a significant rise in vancomycin-resistant *E. faecalis*, which

poses a threat considering that this bacterial species already inherently harbor natural antibiotic-resistant characteristics [36, 39]. Thus, continuous search and discovery of new therapies would hopefully replace those antibiotics to which bacteria may develop resistance. Recently, a new approach has been identified by detecting Anti-QS compounds of herbal origin [40, 41]. Halogenated furanones produced by the benthic marine macroalga *Delisea pulchra* were the first identified anti-QS compounds [42]. Anti-QS compounds are known to have the ability to prohibit bacterial pathogenicity [43]. There is growing interest in implementing novel strategies based on natural compounds to eradicate bacteria, including phytochemicals with anti-biofilm, anti-QS, and anti-oxidant properties that may enhance the efficacy of antibiotics, allowing to reduce their use [44].




Fig. (2). Normalized change values of the *FsrC* gene in *E. faecalis* (ATCC 29,212) and different fractions.




Fig. (3). The effects of different fractions of *S. persica* on the expression of QS gene (*FsrC*) of *E. faecalis* by change fold (average Ct).

S. persica (Miswak-Siwak) is a Middle Eastern native bushy plant whose parts have been used as a natural toothbrush and mouth rinse from time immemorial [30]. Extracts of *S. persica* have even been put into mouthwashes for oral hygiene as it contains trimethylamine, salvadorime chloride, and fluoride in high amounts [45, 46]. *S. persica* is being studied for its antibacterial action against *Streptococcus mutant* and *E. faecalis* [46 - 48]. Other studies reported that *S. persica* has an antibacterial action, allowing it to be used as an irrigant solution in endodontic treatment against endodontic infections, and suggested that it can be used as a root canal irrigant substitute for sodium hypochlorite and chlorhexidine [49, 50]. However, there is currently no published research to determine the exact molecular capability of *S. persica* on QS genes. The present study clearly determined the effects of *S. persica* extracts with different solvents on *E. faecalis* through evaluating the expression of target QS gene (*FsrC*).

In this study, the antibacterial activity of *S. Persica* extract solution was assessed *in vitro* using the broth microdilution method as plant samples are difficult to diffuse in the medium [51, 52]. Previous investigations in Iraq reported that microorganisms in the root canal are highly sensitive to the alcoholic extract of *S. Persica* solution at 15% [49]. The present study also confirms that *E. faecalis* (ATCC 29,212) is highly sensitive to all mentioned fractions of *S. persica*. The antimicrobial effect of *S. Persica* is believed to be due to its high chemical contents of chlorides, tannins, trimethylamine, salvadorine, nitrate, thiocyanate and sulphur [53]. This study confirmed that *S. persica* decreased the expression level of QS gene in *E. faecalis* (ATCC 29,212) and it had antibacterial properties. This finding agrees with previous studies that suggested the Anti-QS ability of *S. persica* against *E. faecalis* [30]. In general, all AQS compounds have a wide range of actions that target practically all bacterial species that depend on QS pathways [10, 11]. This suggests that AQS in *S. persica*

may also function against other opportunistic oral pathogens. Therefore, more studies are required to investigate the AQS ability of *S. persica* against other oral pathogens.

CONCLUSION

The present study concluded that fractions 1, 2, and 3 decreased the expression of the QS gene (*FsrC*), and the highest change fold value (4.99) was recorded by hexane-*S. persica* extracts. Moreover, the lowest concentration of *S. persica* that inhibited the growth of *E. faecalis* was 20mg/ml of fraction 3 (ethyl acetate- *S. persica*). Overall the findings suggest that *S. persica* showed antimicrobial activity against *E. faecalis*.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

HUMAN AND ANIMAL RIGHTS

No humans or animals were used in this study.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

Not applicable.

FUNDING

None

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The corresponding author confirms that ethical considerations were taken for this research, the article is original, and its contents are unpublished. The co-author has read and approved the manuscript for submission.

REFERENCES

- [1] Pinheiro ET, Gomes BP, Ferraz CC, Sousa EL, Teixeira FB, Souza-Filho FJ. Microorganisms from canals of root-filled teeth with periapical lesions. *Int Endod J* 2003; 36(1): 1-11. [http://dx.doi.org/10.1046/j.1365-2591.2003.00603.x] [PMID: 12656508]
- [2] Singh H. Microbiology of endodontic infections. *J Dent Oral Hyg* 2016; 2: 1-4.
- [3] Alghamdi F, Shakir M. The influence of *Enterococcus faecalis* as a dental root canal pathogen on endodontic treatment: A systematic review. *Cureus* 2020; 12(3): e7257. [http://dx.doi.org/10.7759/cureus.7257] [PMID: 32292671]
- [4] Gilmore MS, Lebreton F, van Schaik W. Genomic transition of enterococci from gut commensals to leading causes of multidrug-resistant hospital infection in the antibiotic era. *Curr Opin Microbiol* 2013; 16(1): 10-6. [http://dx.doi.org/10.1016/j.mib.2013.01.006] [PMID: 23395351]
- [5] Chenicheri S, R U, Ramachandran R, Thomas V, Wood A. Insight into oral biofilm: primary, secondary and residual caries and phyto-challenged solutions. *Open Dent J* 2017; 11(1): 312-33. [http://dx.doi.org/10.2174/1874210601711010312] [PMID: 28839480]
- [6] Al-Zubidi M, Widziolk M, Court EK, et al. Identification of novel bacteriophages with therapeutic potential that target *Enterococcus faecalis*. *Infect Immun* 2019; 87(11): e00512-9. [http://dx.doi.org/10.1128/IAI.00512-19] [PMID: 31451618]
- [7] Hastings JW, Greenberg EP. Quorum sensing: the explanation of a curious phenomenon reveals a common characteristic of bacteria. *J Bacteriol* 1999; 181(9): 2667-8. [http://dx.doi.org/10.1128/JB.181.9.2667-2668.1999] [PMID: 10217751]
- [8] Nakayama J, Chen S, Oyama N, et al. Revised model for *Enterococcus faecalis* fsr quorum-sensing system: the small open reading frame fsrD encodes the gelatinase biosynthesis-activating pheromone propeptide corresponding to staphylococcal agrD. *J Bacteriol* 2006; 188(23): 8321-6. [http://dx.doi.org/10.1128/JB.00865-06] [PMID: 16980448]
- [9] Whiteley M, Diggle SP, Greenberg EPJ. Bacterial quorum sensing: the progress and promise of an emerging research area. *Nature* 2017; 551(7680): 313. [http://dx.doi.org/10.1038/nature24624] [PMID: 29144467]
- [10] Bandara HM, Lam OL, Jin LJ, Samaranyake L. Microbial chemical signaling: a current perspective. *Crit Rev Microbiol* 2012; 38(3): 217-49. [http://dx.doi.org/10.3109/1040841X.2011.652065] [PMID: 22300377]
- [11] Giannakara M, Koumandou VLJAM. Evolution of two-component quorum sensing systems. *Access Microbiol* 2022; 4(1): 000303. [http://dx.doi.org/10.1099/acmi.0.000303] [PMID: 35252749]
- [12] Li Y-H, Tian X. Quorum sensing and bacterial social interactions in biofilms. *Sensors (Basel)* 2012; 12(3): 2519-38. [http://dx.doi.org/10.3390/s120302519] [PMID: 22736963]
- [13] Teixeira N d P L. FSR Quorum Sensing: Role in *Enterococcus faecalis* Biology & Host Infection. 2014.
- [14] Nakayama J, Cao Y, Horii T, et al. Gelatinase biosynthesis-activating pheromone: a peptide lactone that mediates a quorum sensing in *Enterococcus faecalis*. *Mol Microbiol* 2001; 41(1): 145-54. [http://dx.doi.org/10.1046/j.1365-2958.2001.02486.x] [PMID: 11454207]
- [15] Teixeira N, Varahan S, Gorman MJ, et al. Drosophila host model reveals new *Enterococcus faecalis* quorum-sensing associated virulence factors. *PLoS One* 2013; 8(5): e64740. [http://dx.doi.org/10.1371/journal.pone.0064740] [PMID: 23734216]
- [16] Hancock LE, Perego M. The *Enterococcus faecalis* fsr two-component system controls biofilm development through production of gelatinase. *J Bacteriol* 2004; 186(17): 5629-39. [http://dx.doi.org/10.1128/JB.186.17.5629-5639.2004] [PMID: 15317767]
- [17] Garsin DA, Sifri CD, Mylonakis E, et al. A simple model host for identifying Gram-positive virulence factors. *Proc Natl Acad Sci USA* 2001; 98(19): 10892-7. [http://dx.doi.org/10.1073/pnas.191378698] [PMID: 11535834]
- [18] Mylonakis E, Engelbert M, Qin X, et al. The *Enterococcus faecalis* fsrB gene, a key component of the fsr quorum-sensing system, is associated with virulence in the rabbit endophthalmitis model. *Infect Immun* 2002; 70(8): 4678-81. [http://dx.doi.org/10.1128/IAI.70.8.4678-4681.2002] [PMID: 12117982]
- [19] Engelbert M, Mylonakis E, Ausubel FM, Calderwood SB, Gilmore MSJ. Contribution of gelatinase, serine protease, and fsr to the pathogenesis of *Enterococcus faecalis* endophthalmitis. *Infect Immun* 2004; 72(6): 3628-33. [http://dx.doi.org/10.1128/IAI.72.6.3628-3633.2004] [PMID: 15155673]
- [20] Nannini EC, Teng F, Singh KV, Murray BE. Decreased virulence of a gls24 mutant of *Enterococcus faecalis* OG1RF in an experimental endocarditis model. *Infect Immun* 2005; 73(11): 7772-4. [http://dx.doi.org/10.1128/IAI.73.11.7772-7774.2005] [PMID: 16239583]
- [21] Denotti G, Piga R, Montaldo C, et al. In Vitro evaluation of *Enterococcus faecalis* adhesion on various endodontic medicaments. *Open Dent J* 2009; 3(1): 120-4. [http://dx.doi.org/10.2174/1874210600903010120] [PMID: 19557151]
- [22] Cook LC, Federle MJF. Peptide pheromone signaling in *Streptococcus* and *Enterococcus*. *Microbiol Rev* 2014; 38(3): 473-92. [http://dx.doi.org/10.1111/1574-6976.12046] [PMID: 24118108]
- [23] Solano C, Echeverez M, Laso I. Biofilm dispersion and quorum sensing. *Curr Opin Microbiol* 2014; 18: 96-104. [http://dx.doi.org/10.1016/j.mib.2014.02.008] [PMID: 24657330]
- [24] Ruhul R, Kataria R. Biofilm patterns in gram-positive and gram-negative bacteria. *Microbiol Res* 2021; 251: 126829. [http://dx.doi.org/10.1016/j.micres.2021.126829] [PMID: 34332222]
- [25] Singh RP, Nakayama J. Development of quorum-sensing inhibitors targeting the fsr system of *Enterococcus faecalis*. *Quorum Sensing vs Quorum Quenching: A Battle with No End in Sight*. Springer 2015; pp. 319-24. [http://dx.doi.org/10.1007/978-81-322-1982-8_25]
- [26] Khalifa L, Gelman D, Shlezinger M, et al. Defeating antibiotic-and phage-resistant *Enterococcus faecalis* using a phage cocktail in vitro and in a clot model. *Front Microbiol* 2018; 9: 326. [http://dx.doi.org/10.3389/fmicb.2018.00326] [PMID: 29541067]
- [27] Sanchez Armengol E, Harmanci M, Laffleur F. Current strategies to determine antifungal and antimicrobial activity of natural compounds. *Microbiol Res* 2021; 252: 126867. [http://dx.doi.org/10.1016/j.micres.2021.126867] [PMID: 34521051]
- [28] Jose PA, Maharshi A, Jha B. Actinobacteria in natural products research: Progress and prospects. *Microbiol Res* 2021; 246: 126708. [http://dx.doi.org/10.1016/j.micres.2021.126708] [PMID: 33529791]
- [29] Williams AN, Stavrinides J. Pantoea Natural Product 3 is encoded by an eight-gene biosynthetic gene cluster and exhibits antimicrobial activity against multi-drug resistant *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. *Microbiol Res* 2020; 234: 126412. [http://dx.doi.org/10.1016/j.micres.2020.126412] [PMID: 32062363]
- [30] Rezaei A, Oyong GG, Borja VB, et al. Molecular screening of anti-quorum sensing capability of *Salvadora persica* on *Enterococcus faecalis*. *J Hard Tissue Biol* 2011; 20(2): 115-24. [http://dx.doi.org/10.2485/jhtb.20.115]
- [31] Balto H, Al-Sanie I, Al-Beshri S, Aldrees A. Effectiveness of *Salvadora persica* extracts against common oral pathogens. *Saudi Dent J* 2017; 29(1): 1-6. [http://dx.doi.org/10.1016/j.sdentj.2016.11.001] [PMID: 28270703]
- [32] Arias-Moliz MT, Ferrer-Luque CM, González-Rodríguez MP, Navarro-Escobar E, de Freitas MF, Baca P. Antimicrobial activity and *Enterococcus faecalis* biofilm formation on chlorhexidine varnishes. *Med Oral Patol Oral Cir Bucal* 2012; 17(4): e705-9. [http://dx.doi.org/10.4317/medoral.17680] [PMID: 22322495]
- [33] Harborne A. Phytochemical methods a guide to modern techniques of plant analysis. Springer Science & Business Media 1998.
- [34] Eloff JNJ. A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Med* 1998; 64(8): 711-3. [http://dx.doi.org/10.1055/s-2006-957563] [PMID: 9933989]
- [35] Wayne P. Clinical and laboratory standards institute. Performance standards for antimicrobial susceptibility testing. CLSI document 2011.
- [36] Shepard BD, Gilmore MS. Differential expression of virulence-related

- genes in *Enterococcus faecalis* in response to biological cues in serum and urine. *Infect Immun* 2002; 70(8): 4344-52. [http://dx.doi.org/10.1128/IAI.70.8.4344-4352.2002] [PMID: 12117944]
- [37] Herdiyati Y, Atmaja HE, Satari MH, Kurnia D. Potential antibacterial flavonoid from buah merah (Lam.) against pathogenic oral bacteria of ATCC 29212. *Open Dent J* 2020; 14(1) [http://dx.doi.org/10.2174/1874210602014010433]
- [38] Qin X, Singh KV, Weinstock GM, Murray BE. Characterization of *fsr*, a regulator controlling expression of gelatinase and serine protease in *Enterococcus faecalis* OG1RF. *J Bacteriol* 2001; 183(11): 3372-82. [http://dx.doi.org/10.1128/JB.183.11.3372-3382.2001] [PMID: 11344145]
- [39] Borzini L, Condò R, De Dominicis P, Casaglia A, Cerroni L. Root canal irrigation: Chemical agents and plant extracts against *Enterococcus faecalis*. *Open Dent J* 2016; 10(1): 692-703. [http://dx.doi.org/10.2174/1874210601610010692] [PMID: 28217184]
- [40] Ni N, Li M, Wang J, Wang B. Inhibitors and antagonists of bacterial quorum sensing. *Med Res Rev* 2009; 29(1): 65-124. [http://dx.doi.org/10.1002/med.20145] [PMID: 18956421]
- [41] Anand S, Rajan M, Venkateshabu N, Kandaswamy D, Shrivaya Y, Rajeswari K. Evaluation of the Antibacterial Efficacy of *Azadirachta Indica*, *Commiphora Myrrha*, *Glycyrrhiza Glabra* Against *Enterococcus faecalis* using Real Time PCR. *Open Dent J* 2016; 10(1): 160-5. [http://dx.doi.org/10.2174/1874210601610010160] [PMID: 27386000]
- [42] Aldawsari MF, Khafagy E-S, Saqr AA, et al. Tackling virulence of *Pseudomonas aeruginosa* by the natural furanone sotolon. *Antibiotics (Basel)* 2021; 10(7): 871. [http://dx.doi.org/10.3390/antibiotics10070871] [PMID: 34356792]
- [43] Asfour HZ. Anti-quorum sensing natural compounds. *J Microsc Ultrastruct* 2018; 6(1): 1-10. [http://dx.doi.org/10.4103/JMAU.JMAU_10_18] [PMID: 30023261]
- [44] Alibi S, Ben Selma W, Ramos-Vivas J, et al. Anti-oxidant, antibacterial, anti-biofilm, and anti-quorum sensing activities of four essential oils against multidrug-resistant bacterial clinical isolates. *Curr Res Transl Med* 2020; 68(2): 59-66. [http://dx.doi.org/10.1016/j.retram.2020.01.001] [PMID: 32192922]
- [45] Al-Bayaty FH, Al-Koubaisi, A. H.; Ali, N. A. W.; Abdulla, M. A., Effect of mouth wash extracted from *Salvadora persica* (Miswak) on dental plaque formation: A clinical trail. *J Med Plants Res* 2013; 4(14): 1446-58.
- [46] Almas K J. The antimicrobial effects of extracts of *Azadirachta indica* (Neem) and *Salvadora persica* (Arak) chewing sticks. *Indian Soc Dent Res* 1999; 10(1): 23-6.
- [47] Balhaddad AA, Mokeem L, Melo MAS, Gregory RL. Antibacterial activities of methanol and aqueous extracts of *Salvadora persica* against *Streptococcus mutans* biofilms: An *in vitro* study. *Dent J* 2021; 9(12): 143. [http://dx.doi.org/10.3390/dj9120143] [PMID: 34940040]
- [48] Khalessi AM, Pack AR, Thomson WM, Tompkins GR. An *in vivo* study of the plaque control efficacy of Persica: A commercially available herbal mouthwash containing extracts of *Salvadora persica*. *Int Dent J* 2004; 54(5): 279-83. [http://dx.doi.org/10.1111/j.1875-595X.2004.tb00294.x] [PMID: 15503853]
- [49] Al-Sabawi N, Al Sheikh Abdal A, Taha M Y. The antimicrobial activity of *Salvadora persica* solution (miswak-siwak) as root canal irrigant (a comparative study). *J Pure Appl Sci* 2007; 4(3): 69-91.
- [50] Almas K. The effect of *Salvadora persica* extract (miswak) and chlorhexidine gluconate on human dentin: A SEM study. *J Contemp Dent Pract* 2002; 3(3): 27-35. [http://dx.doi.org/10.5005/jcdp-3-3-27] [PMID: 12239575]
- [51] Verpoorte R. Antimicrobially active alkaloids. *Springer Alkaloids* 1998; pp. 397-433. [http://dx.doi.org/10.1007/978-1-4757-2905-4_17]
- [52] Estrela CR, Estrela C, Reis C, Bammann LL, Pécora JD. Control of microorganisms *in vitro* by endodontic irrigants. *Braz Dent J* 2003; 14(3): 187-92. [http://dx.doi.org/10.1590/S0103-64402003000300009] [PMID: 15057395]
- [53] Akhtar MS, Ajmal M. Significance of chewing-sticks (miswaks) in oral hygiene from a pharmacological view-point. *J Pak Med Assoc* 1981; 31(4): 89-95. [PMID: 6785501]